US ISSN 0046 225X

ENVIRONMENTAL ENTOMOLOGY

Supplement to Environmental Entomology Vol. 30 No. 6 December 2001



Published by the Entomological Society of America

Volume 30 Number 6

December 2001

Environmental Entomology

Volume 30
Number 6
December 2001

JESSE A. LOGAN
DAVID W. ONSTAD
EDWARD L. VARGO
THOMAS W. SAPPINGTON
Co-Editors

ABELARDO C. MONCAYO
Book & Media Review Editor

ALAN KAHAN
Director of Communications

MAGGIE A. MEITZLER
Managing Editor/ Manuscript Editor

JEANNIE WIMBUSH ALLEN
Production Editor

EDITORIAL BOARD

P. H. ADLER, *Chair*
Clemson, SC
Section A, 2002

M. W. BROWN
Kearneysville, WV
Section C, 2002

R. A. WEINZIERL
Urbana, IL
Section E

R. E. LEE
Oxford, OH
Section B, 2004

S. PAULSON
Blacksburg, VA
Section D, 2003

M. J. WEISS
Moscow, ID
Section F, 2001

GOVERNING BOARD

L. L. LARSON
Indianapolis, IN
President

K. S. SHIELDS
Hamden, CT
Secretary-Treasurer

F. E. GILSTRAP
College Station, TX
Southwestern Branch

G. R. MULLEN
Auburn, AL
Section D

J. E. McPHERSON
Carbondale, IL
Vice President

B. A. NAULT
Painter,VA
Eastern Branch

M. A. IVIE
Bozeman, MT
Section A

W. F. GIMPEL
Annapolis, MD
Section E

S. S. QUISENBERRY
Bozeman, MT
Past-President

S. H. HUTCHINS
Indianapolis, IN
North Central Branch

D. B. GELMAN
Beltsville, MD
Section B

S. T. KAMBLE
Lincoln, NE
Section F

Z. B. MAYO, JR.
Lincoln, NE
Vice President-Elect

R. A. REDAK
Riverside, CA
Pacific Branch

L. A. LACEY
Wapato, WA
Section C

P. G. LETTICE
Lanham, MD
Executive Director

© 2001, Entomological Society of America

Environmental Entomology (ISSN 0013-8746) is published bimonthly (February, April, June, August, October, and December) by the Entomological Society of America, 9301 Annapolis Road, Lanham, MD 20706–3115; (301) 731-4535; FAX (301) 731-4538. The journal is printed by Cadmus Journal Services, 2905 Byrdhill Road, Richmond, VA 23228. Periodicals postage is paid at Lanham, MD, and at additional mailing offices.

Postmaster: Send change of address forms to *Environmental Entomology*, ESA, 9301 Annapolis Road, Lanham, MD 20706–3115.

Inquiries regarding content, subscriptions, advertising, availability of back issues, change of address, general business matters, and manuscripts should be sent to the Publications Office, ESA, 9301 Annapolis Road, Lanham, MD 20706–3115. For information about the preparation of manuscripts, authors are urged to consult "Information for Contributors," which is printed on the inside back cover of most issues, or our *Publishing with ESA Style Guide*, located on the ESA website at http://www.entsoc.org.

Articles report the results of research only. Mention of a pesticide or a commercial or proprietary product does not constitute an endorsement or recommendation for its use. Registered names and trademarks, etc., used in this publication, even without specific indication thereof, are not to be considered unprotected by law. All articles and items published in the Journal are signed and, unless otherwise stated, reflect the individual views of the authors and not necessarily the official points of view held by ESA. ESA assumes no responsibility for the completeness of the articles or other items or the accuracy of the conclusions reached therein.

Photocopies. Permission to photocopy journal articles for internal or personal use is granted by the Entomological Society of America provided that a fee of $2.00 per copy is paid directly to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923. Individuals may make a single copy for personal use without charge. This consent does not extend to other kinds of copying, such as that for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale.

Subscriptions. 2001 prices: nonmember individuals, $95; institutions, $191; nonmember individual-online $95; and institutions-online $191. Also, add $20 for shipping outside the continental United States or $36 for airmail. Send payment to EE Subscriptions, ESA, P.O. Box 177, Hyattsville, MD 20781. Inquiries should be directed to ESA Sales, 9301 Annapolis Road, Lanham, MD 20706–3115; (301) 731-4535; FAX (301) 731-4538.

Back Issues	ESA member domestic	ESA member foreign	Non-member domestic	Non-member foreign
Single Issue	$10.40	$11.40	$34.00	$35.00
Volume	62.40	68.40	204.00	210.00

All prices include handling and shipping by surface mail. For a list of available issues, write to ESA Sales, 9301 Annapolis Road, Lanham, MD 20706–3115. Back volumes are available in microform from University Microfilms International, 300 North Zeeb Road, Ann Arbor, MI 48106.

CORRECTED CUMULATIVE SUBJECT INDEX
Environmental Entomology
Volume 30, 2001

A

abundance,
dispersal, 267
habitats, 267
highways, 267
populations, 267
Solenopsis invicta, 267
accumulated air mean subfreezing temperature,
bean leaf beetle, 562
Cerotoma trifurcata, 562
multiple regression stepwise selection procedure analysis, 562
population dynamics, 562
acetic acid,
attractant, 656
Lacanobia subjuncta, 656
Mamestra configurata, 656
trapping, 656
Xestia c-nigrum, 656
acetylcholinesterase,
bark beetles, 803
carbaryl, 803
environmental safety, 803
Pinaceae, 803
Scolytidae, 803
Acleris gloverana,
age-specific survival ratios, 883
dispersal, 883
Douglas-fir tussock moth, 883
population dynamics, 883
western spruce budworm, 883
Acrolepiopsis assectella,
behavior, 288
emigration, 288
intercrop, 288
oviposition, 288
Plutella xylostella, 288
Acyrthosiphon pisum,
Coccinellidae, 964
Coleomegilla maculata, 964
fatty acids, 964
Hippodamia convergens, 964
tritrophic interactions, 964
Adelgidae,
diapause, 877
photoperiod, 877
temperature, 877
Tsuga spp. aestivation, 877
adult longevity,
pheromone trap catch, 999
pupal weight, 999
Rhyacionia frustrana, 999
adult size,
aphid prey, 1182
coccinellid predator, 1182
functional response, 1182
larval food supply, 1182
Menochilus sexmaculatus, 1182
Aedes aegypti,
Aedes albopictus, 855
global warming, 855
larval development, 855
seasonal environment, 855
survival, 855
Aedes albopictus,
Aedes aegypti, 855
global warming, 855
larval development, 855
seasonal environment, 855
survival, 855
Aedes sollicitans,
control, 388
mosquito, 388
starch, 388
xanthan gum, 388
aerial take-off,
diffusion coefficient, 1112
infochemicals, 1112
spider mites, 1112
walking speed, 1112
aflatoxins,
mycotoxins, 817
protein metabolism, 817
Spodoptera littoralis, 817
Agapeta zoegana,
biological control, 589
Centaurea maculosa, 589
Cyphocleonus achates, 589
larval sampling, 589
spotted knapweed, 589
age-specific survival ratios,
Acleris gloverana, 883
dispersal, 883
Douglas-fir tussock moth, 883
population dynamics, 883
western spruce budworm, 883
aggregation,
ambush predator, 51
chemical defense, 51
frogs, 51
warning coloration, 51
aggregation sites,
dispersal, 700
mate seeking, 700
Ostrinia nubilalis, 700
refuge, 700
surface airflow, 700
Agraulis vanillae,
Eurema lisa, 1052
Junonia coenia, 1052
phenology, 1052
Phoebis sennae, 1052
Urbanus proteus, 1052
agroecosystems,
boundaries, 225
dispersal, 225
multivariate analysis, 225
predators, 225
Triticum aestivum, 225
Aiolomorphus rhopaloides,
bamboo, 1098
Diomorus aiolomorphi, 1098
gall maker, 1098
phenology, 1098
air pollution,
Colorado potato beetle, 824
host stress, 824
Leptinotarsa decemlineata, 824
ozone, 824
potato, 824
alatae,
apterae, 1047
density, 1047
development, 1047
Toxoptera citricida, 1047
Aleurothrixus aepim,
ant-homopteran interactions, 295
Croton floribundus, 295
insect–plant interactions, 295
mutualism, 295
sooty mold, 295
alfalfa looper,
attractant, 667
Autographa californica, 667
phenylacetaldehyde, 667
trap, 667
Allorhogas pyralophagus,
Eoreuma loftini, 129
functional response, 129
reproduction, 129
almond,
dormant spray, 70
esfenvalerate, 70
Galendromus occidentalis, 70
permethrin, 70
residue, 70
Alphitobius diaperinus,
broiler production, 556
geographic information system, 556
lesser mealworm, 556

spatial model, 536
alternating current electrical penetration graph system,
corn leaf aphid, 1041
plant resistance, 1041
plant water stress, 1041
probing behavior, 1041
Russian wheat aphid, 1041
Amblyomma americanum,
distribution, 568
habitats, 568
Ixodes scapularis, 568
ticks, 568
ambush predator,
aggregation, 51
chemical defense, 51
frogs, 51
warning coloration, 51
ambushers,
diversity, 1021
hunting spiders, 1021
predatory arthropods, 1021
sampling methods, 1021
web spiders, 1021
Anoplolepis longipes,
biological control, 1028
community structure, 1028
Nezara viridula, 1028
Pheidole megacephala, 1028
ant–homopteran interaction,
Formica, 757
leafhoppers, 757
mutualism, 757
myrmecophile, 757
ant-homopteran interactions,
Aleurothrixus aepim, 295
Croton floribundus, 295
insect–plant interactions, 295
mutualism, 295
sooty mold, 295
Anthonomus grandis,
cotton pest management, 550
diapause, 550
overwintering survivorship, 550
Anticarsia gemmatalis, 76, 1164
beneficial nontarget organisms, 1164
biological control, 1164
biopesticide, 1164
cultural control, 76
environmental risk assessment, 1164
Glycine max, 76
Podisus nigrispinus, 1164
Pseudoplusia includens, 76
Spissistilus festinus, 76
stink bugs, 76
antipredator behavior,
Diabrotica undecimpunctata howardi, 327
predation, 327
sexual differences, 327
trophic cascade, 327
ants, 425, 842
Cyclocephala lurida, 425
forest, 842
Formicidae, 842
habitat, 842
Popillia japonica, 425
predators, 425
Solenopsis molesta, 425
turfgrass, 425
Aphantorhaphopsis samarensis,
biological control, 605
host range, 605
host specificity, 605
host suitability, 605
Aphelinus albipodus,
Aphelinus asychis, 578
aphid parasitoids, 578
biological control, 578
Diaeretiella rapae, 578
Russian wheat aphid, 578
Aphelinus asychis,
Aphelinus albipodus, 578
aphid parasitoids, 578
biological control, 578
Diaeretiella rapae, 578
Russian wheat aphid, 578
aphid parasitoids,
Aphelinus albipodus, 578
Aphelinus asychis, 578
biological control, 578
Diaeretiella rapae, 578
Russian wheat aphid, 578
aphid prey,
adult size, 1182
coccinellid predator, 1182
functional response, 1182
larval food supply, 1182
Menochilus sexmaculatus, 1182
Aphididae,
habitat selection, 371
migratory flight, 371
suction trap, 371
aphids,
corn, 1127
corn aphids, 1127
development, 1127
fecundity, 1127
temperatures, 1127
Aphis gossypii,
biological control, 785
Coccinellidae, 785
functional response, 785
population dynamics, 785
Scymnus creperus, 785
apple, 415, 763
biodiversity, 415
biological control, 415
Cydia pomonella, 763
diel periodicity, 415
fruit damage, 763
inherited sterility, 763
orchard management, 415
apterae,
alatae, 1047
density, 1047
development, 1047
Toxoptera citricida, 1047
aquatic barrier trap,
aquatic intercept trap, 770
Lissorhoptrus oryzophilus, 770
rice integrated pest management, 770
rice water weevil, 770
weevil trapping, 770
aquatic intercept trap,
aquatic barrier trap, 770
Lissorhoptrus oryzophilus, 770
rice integrated pest management, 770
rice water weevil, 770
weevil trapping, 770
Argentine ant,
biological invasion, 742
host specificity, 742
Linepithema humile, 742
Phoridae, 742
Pseudacteon, 742
arthropods, 711, 1017
cover crops, 1017
marking, 711
movement, 711
predator, 711
rubidium, 711
Solenopsis invicta, 1017
sugarcane, 1017
Asclepias,
herbivory, 333
subterranean herbivory, 333
Tetraopes, 333
Asian longhorned beetle,
Cerambycidae, 1036
diffusion, 1036
invasive species, 1036
mark-recapture, 1036
Aspen-FACE,
Betula papyrifera, 1119
CO_2, 1119
O_3, 1119
Orgyia leucostigma whitemarked tussock moth, 1119
Asteraceae,
biological control, 439
Centaurea solstitialis, 439
Curculionidae, 439
Eustenopus villosus, 439
phenological synchrony, 439
attractant, 656, 667
acetic acid, 656
alfalfa looper, 667
Autographa californica, 667
Lacanobia subjuncta, 656

Mamestra configurata, 656
phenylacetaldehyde, 667
trap, 667
trapping, 656
Xestia c-nigrum, 656
Attractants,
food volatiles, 157
isobutanol, 157
Vespidae, 157
Vespula germanica, 157
yellowjackets, 157
Autographa californica,
alfalfa looper, 667
attractant, 667
phenylacetaldehyde, 667
trap, 667
avocado,
scarring on avocado fruit, 932
Scirtothrips perseae, 932
thresholds, 932
thrips numbers, 932

B

Bacillus thuringiensis, 409, 495, 625, 642, 728, 1174
Bt-corn, 495
Compsilura, 1174
corn, 625, 728
Cry1F, 642
Danaus plexippus, 495
degradation, 642
insect resistance management, 409
insidious flower bug, 625
larval survival, 728
Lymantria dispar, 1174
monarch butterfly, 495
nontarget effects, 495
Orius insidiosus, 625
Ostrinia nubilalis, 409, 728
pollen, 409
predator, 625
resistance management, 728
soil, 642
sublethal effects, 1174
transgenic, 495
transgenic corn, 409
weeds, 728
Zea mays, 409
Bacillus thuringiensis tenebrionis,
Colorado potato beetle, 524
plant–insect interactions, 524
potato, 524
potato aphid, 524
transgenic classical resistance, 524
bacteria,
bio-control, 443
Coptotermes formosanus, 443
fungus, 443
Serratia marcescens, 443
termite, 443
bamboo,
Aiolomorphus rhopaloides, 1098
Diomorus aiolomorphi, 1098
gall maker, 1098
phenology, 1098
bark beetles, 215, 803, 993
acetylcholinesterase, 803
carbaryl, 803
Dendroctonus pseudotsugae, 215
disturbance, 215
Douglas-fir beetle, 215
Enoclerus sphegeus, 993
environmental safety, 803
hazard rating model, 215
kairomone, 993
Pinaceae, 803
Pityogenes carinulatus, 993
risk classification model, 215
Scolytidae, 803
Temnochila chlorodia, 993
Thanasimus undatulus, 993
bean leaf beetle,
accumulated air mean subfreezing temperature, 562
Cerotoma trifurcata, 562
multiple regression stepwise selection procedure analysis, 562
population dynamics, 562
beet armyworm,
cotton, 511
egg mass, 511
oviposition, 511
pigweed, 511
Spodoptera exigua, 511
beetles,
deciduous forests, 484
disturbance, 484
forest floor, 484
richness, 484
Shannon diversity index, 484
behavior, 288, 466, 1012
Acrolepiopsis assectella, 288
biological control, 1012
emigration, 288
Hippodamia convergens, 1012
intercrop, 288
mole crickets, 466
Myzus nicotianae, 1012
olfactory cues, 1012
oviposition, 288, 466
Plutella xylostella, 288
prey searching, 1012
Scapteriscus borellii, 466
soil moisture, 466
Bemisia tabaci, 55, 89, 720
biotype B, 720
biotype Q, 720
cantaloupe, 55
cotton, 55
development, 55, 720
fecundity, 55
intercropping, 89
pepper, 720
polyculture, 89
survival, 55
Trialeurodes vaporariorum, 89
vector management, 89
beneficial nontarget organisms,
Anticarsia gemmatalis, 1164
biological control, 1164
biopesticide, 1164
environmental risk assessment, 1164
Podisus nigrispinus, 1164
Benin,
environmental variables, 1135
flight activity, 1135
modeling, 1135
pheromone trapping, 1135
Prostephanus truncatus, 1135
Betula papyrifera,
Aspen-FACE, 1119
CO_2, 1119
O_3, 1119
Orgyia leucostigma whitemarked tussock moth, 1119
bio-control,
bacteria, 443
Coptotermes formosanus, 443
fungus, 443
Serratia marcescens, 443
termite, 443
biocontrol,
Coptotermes formosanus, 449
eicosanoid, 449
immunity, 449
insect pathology, 449
Serratia marcescens, 449
biodiversity,
apple, 415
biological control, 415
diel periodicity, 415
orchard management, 415
biogeographical range,
distribution, 645
modeling, 645
quarantine, 645
Solenopsis invicta, 645
United States, 645
biological control, 147, 415, 439, 578, 589, 605, 612, 617, 637, 785, 982, 1012, 1028, 1164
Agapeta zoegana, 589
Anoplolepis longipes, 1028
Anticarsia gemmatalis, 1164
Aphantorhaphopsis samarensis, 605
Aphelinus albipodus, 578
Aphelinus asychis, 578

aphid parasitoids, 578
Aphis gossypii, 785
apple, 415
Asteraceae, 439
behavior, 1012
beneficial nontarget organisms, 1164
biodiversity, 415
biopesticide, 1164
Carduus nutans, 612
Centaurea maculosa, 589
Centaurea solstitialis, 439
Coccinellidae, 785
community structure, 1028
Curculionidae, 439
Cyphocleonus achates, 589
development, 637
Diaeretiella rapae, 578
diel periodicity, 415
environmental risk assessment, 1164
Eustenopus villosus, 439
functional response, 785
Hippodamia convergens, 1012
host range, 605
host specificity, 605
host suitability, 605
larval sampling, 589
life history, 147, 982
Lysiphlebus testaceipes, 637
Musca domestica, 147, 982
Muscidifurax raptorellus, 982
Muscidifurax zaraptor, 147
musk thistle, 612
Myzus nicotianae, 1012
Nezara viridula, 1028
olfactory cues, 1012
orchard management, 415
Oryzaephilus surinamensis, 617
parasitoid, 637
Pheidole megacephala, 1028
phenological synchrony, 439
Plodia interpunctella, 617
Podisus nigrispinus, 1164
population dynamics, 785
prey searching, 1012
red imported fire ant, 612
Rhinocyllus conicus, 612
Rhyzopertha dominica, 617
Russian wheat aphid, 578
Schizaphis graminum, 637
Scymnus creperus, 785
Solenopsis invicta, 612
spotted knapweed, 589
Stomoxys calcitrans, 147, 982
stored products, 617
temperature, 147, 982
Tribolium castaneum, 617
biological control of weeds,
Carthamus tinctorius, 953
Centaurea solstitialis, 953
choice tests, 953
nontarget impact, 953
realized host range, 953
biological invasion,
Argentine ant, 742
host specificity, 742
Linepithema humile, 742
Phoridae, 742
Pseudacteon, 742
biopesticide,
Anticarsia gemmatalis, 1164
beneficial nontarget organisms, 1164
biological control, 1164
environmental risk assessment, 1164
Podisus nigrispinus, 1164
biotype B,
Bemisia tabaci, 720
biotype Q, 720
development, 720
pepper, 720
biotype Q,
Bemisia tabaci, 720
biotype B, 720
development, 720
pepper, 720
black cutworm moth,
calling, 189
migration, 189
periodicity, 189
reproductive diapause, 189
sex pheromone, 189
blueberry maggot fly,
emergence models, 925
highbush blueberries, 925
postdiapause development, 925
Rhagoletis mendax, 925
body water content,
cuticular permeability, 631
Dryas iulia, 631
Heliconius, 631
surface area, 631
boll weevil,
cultural management, 64
overwintering, 64
pheromone synergist, 64
trapping, 64
bollworm, 112, 431
cotton, 112, 431
enzyme-linked immunosorbent assay, 431
life table, 112
Orius, 431
predation, 431
serological assay, 431
bone pile,
foraging, 123
natural enemies, 123
predator, 123
red imported fire ant, 123
refuse pile, 123
boundaries,
agroecosystems, 225
dispersal, 225
multivariate analysis, 225
predators, 225
Triticum aestivum, 225
Brachycera,
Microtrombidiidae, 903
Parasitization, 903
Tachinidae, 903
Therevidae, 903
Brassicaceae,
lady beetle, 947
Myzus persicae, 947
tritrophic interactions, 947
broiler production,
Alphitobius diaperinus, 556
geographic information system, 556
lesser mealworm, 556
spatial model, 556
Bt-corn,
Bacillus thuringiensis, 495
Danaus plexippus, 495
monarch butterfly, 495
nontarget effects, 495
transgenic, 495

C

Coptotermes formosanus,
Reticulitermes flavipes, 545
termite tunneling, 545
tunneling geometry, 545
calling,
black cutworm moth, 189
migration, 189
periodicity, 189
reproductive diapause, 189
sex pheromone, 189
Callirhytis cornigera,
Cynipidae, 280
density-dependent parasitism, 280
galls, 280
spatial distribution, 280
calorespirometry,
convergent lady beetle, 811
Hippodamia convergens, 811
isothermal calorimetry, 811
respiration, 811
temperature response, 811
cantaloupe,
Bemisia tabaci, 55
cotton, 55
development, 55
fecundity, 55
survival, 55
carbaryl,
acetylcholinesterase, 803
bark beetles, 803
environmental safety, 803

Pinaceae, 803
Scolytidae, 803
carbon-nutrient balance,
ectomycorrhiza, 849
gypsy moth, 849
herbivore-plant interactions, 849
Pisolithus tinctorius, 849
Quercus, 849
Carduus nutans,
biological control, 612
musk thistle, 612
red imported fire ant, 612
Rhinocyllus conicus, 612
Solenopsis invicta, 612
Caribbean region,
invasive pests, 339
Maconellicoccus hirsutus, 339
quarantine, 339
sex pheromones, 339
Carposina sasakii,
development, 298
stage emergence models, 298
Carthamus tinctorius,
biological control of weeds, 953
Centaurea solstitialis, 953
choice tests, 953
nontarget impact, 953
realized host range, 953
Centaurea maculosa,
Agapeta zoegana, 589
biological control, 589
Cyphocleonus achates, 589
larval sampling, 589
spotted knapweed, 589
Centaurea solstitialis, 439, 953
Asteraceae, 439
biological control, 439
biological control of weeds, 953
Carthamus tinctorius, 953
choice tests, 953
Curculionidae, 439
Eustenopus villosus, 439
nontarget impact, 953
phenological synchrony, 439
realized host range, 953
Cerambycidae, 17, 1036
Asian longhorned beetle, 1036
diffusion, 1036
intraguild predation, 17
invasive species, 1036
Ips, 17
Ips calligraphus, 17
mark-recapture, 1036
Monochamus carolinensis, 17
Scolytidae, 17
Cerotoma trifurcata,
accumulated air mean subfreezing temperature, 562
bean leaf beetle, 562
multiple regression stepwise selection procedure analysis, 562
population dynamics, 562
chemical communication,
Coptotermes formosanus, 457
Reticulitermes flavipes, 457
trail pheromone, 457
chemical defense,
aggregation, 51
ambush predator, 51
frogs, 51
warning coloration, 51
choice tests,
biological control of weeds, 953
Carthamus tinctorius, 953
Centaurea solstitialis, 953
nontarget impact, 953
realized host range, 953
Choristoneura occidentalis,
mortality, 136
multicapsid nucleopolyhedrovirus, 136
nucleopolyhedrovirus, 136
sublethal effect, 136
Chrysomela scripta,
larval performance, 748
Populus, 748
short-rotation forestry, 748
Chrysoperla plorabunda,
extrafloral nectar, 595
generalist predator, 595
omnivory, 595
clearcut,
forest management, 400
insect sampling, 400
selection, 400
shelterwood, 400
vertical distribution of insect fauna, 400
CO_2,
Aspen-FACE, 1119
Betula papyrifera, 1119
O_3, 1119
Orgyia leucostigma whitemarked tussock moth, 1119
coccinellid predator,
adult size, 1182
aphid prey, 1182
functional response, 1182
larval food supply, 1182
Menochilus sexmaculatus, 1182
Coccinellidae, 785, 964
Acyrthosiphon pisum, 964
Aphis gossypii, 785
biological control, 785
Coleomegilla maculata, 964
fatty acids, 964
functional response, 785
Hippodamia convergens, 964
population dynamics, 785
Scymnus creperus, 785
tritrophic interactions, 964
cold hardiness,
invading species, 181
range limit, 181
southwestern corn borer, 181
winter mortality, 181
Coleomegilla maculata,
Acyrthosiphon pisum, 964
Coccinellidae, 964
fatty acids, 964
Hippodamia convergens, 964
tritrophic interactions, 964
Coleoptera,
Lagria hirta, 686
Lagriidae, 686
life-history trait correlation, 686
reproduction, 686
Coleoptera: Silvanidae population regulation,
density dependence, 1061
habitat deterioration, 1061
Oryzaephilus surinamensis, 1061
colonization,
flight, 306
impact traps, 306
Leptinotarsa decemlineata, 306
plastic-lined trenches, 306
walking, 306
Colorado potato beetle, 350, 524, 824
air pollution, 824
Bacillus thuringiensis tenebrionis, 524
geostatistics, 350
global positioning systems, 350
host stress, 824
insect spatial dynamics, 350
Leptinotarsa decemlineata, 824
ozone, 824
plant–insect interactions, 524
population dynamics, 350
potato, 524, 824
potato aphid, 524
precision agriculture, 350
transgenic classical resistance, 524
colored sticky traps,
Eucalyptus globulus, 365
heteroblastic host plant, 365
Psyllid, 365
community structure,
Anoplolepis longipes, 1028
biological control, 1028
Nezara viridula, 1028
Pheidole megacephala, 1028

competition,
in-host compatibility, 1158
Lygus lineolaris, 1158
parasitism, 1158
Peristenus, 1158
tarnished plant bug, 1158
component ratio,
pheromone quantity, 1006
Rhyacionia frustrana, 1006
sex pheromone, 1006
synthetic bait, 1006
Compsilura,
Bacillus thuringiensis, 1174
Lymantria dispar, 1174
sublethal effects, 1174
control,
Aedes sollicitans, 388
mosquito, 388
starch, 388
xanthan gum, 388
convergent lady beetle,
calorespirometry, 811
Hippodamia convergens, 811
isothermal calorimetry, 811
respiration, 811
temperature response, 811
Copidosoma floridanum,
diflubenzuron, 794
polyembryony, 794
soybean, 794
Coptotermes formosanus, 262, 443, 449, 457
bacteria, 443
bio-control, 443
biocontrol, 449
chemical communication, 457
DNA fingerprinting, 262
eicosanoid, 449
fungus, 443
genetic structure, 262
immunity, 449
insect pathology, 449
Reticulitermes flavipes, 457
Serratia marcescens, 443, 449
termite, 443
trail pheromone, 457
corn, 625, 728, 1127
aphids, 1127
Bacillus thuringiensis, 625, 728
corn aphids, 1127
development, 1127
fecundity, 1127
insidious flower bug, 625
larval survival, 728
Orius insidiosus, 625
Ostrinia nubilalis, 728
predator, 625
resistance management, 728
temperatures, 1127
weeds, 728
corn aphids,
aphids, 1127
corn, 1127
development, 1127
fecundity, 1127
temperatures, 1127
corn leaf aphid,
alternating current electrical penetration graph system, 1041
plant resistance, 1041
plant water stress, 1041
probing behavior, 1041
Russian wheat aphid, 1041
Cotesia flavipes,
parasitoid, 869
spatial interpolation, 869
spatial-temporal population dynamics, 869
validation, 869
cotton, 55, 112, 431, 511
beet armyworm, 511
Bemisia tabaci, 55
bollworm, 112, 431
cantaloupe, 55
development, 55
egg mass, 511
enzyme-linked immunosorbent assay, 431
fecundity, 55
life table, 112
Orius, 431
oviposition, 511
pigweed, 511
predation, 431
serological assay, 431
Spodoptera exigua, 511
survival, 55
cotton pest management,
Anthonomus grandis, 550
diapause, 550
overwintering survivorship, 550
cover crops,
arthropods, 1017
Solenopsis invicta, 1017
sugarcane, 1017
Croton floribundus,
Aleurothrixus aepim, 295
ant-homopteran interactions, 295
insect–plant interactions, 295
mutualism, 295
sooty mold, 295
Cry1F,
Bacillus thuringiensis, 642
degradation, 642
soil, 642
cultural control, 76, 204
Anticarsia gemmatalis, 76
Empoasca fabae, 204
Glycine max, 76
intercropping, 204
Pseudoplusia includens, 76
resource concentration hypothesis, 204
soybean, 204
Spissistilus festinus, 76
stink bugs, 76
wheat, 204
cultural management,
boll weevil, 64
overwintering, 64
pheromone synergist, 64
trapping, 64
Curculionidae,
Asteraceae, 439
biological control, 439
Centaurea solstitialis, 439
Eustenopus villosus, 439
phenological synchrony, 439
cuticular permeability,
body water content, 631
Dryas iulia, 631
Heliconius, 631
surface area, 631
Cyclocephala borealis,
Cyclocephala hirta, 899
Cyclocephala longula, 899
Cyclocephala lurida, 899
Cyclocephala pasadenae, 899
masked chafer beetles, 899
Cyclocephala hirta,
Cyclocephala borealis, 899
Cyclocephala longula, 899
Cyclocephala lurida, 899
Cyclocephala pasadenae, 899
masked chafer beetles, 899
Cyclocephala longula,
Cyclocephala borealis, 899
Cyclocephala hirta, 899
Cyclocephala lurida, 899
Cyclocephala pasadenae, 899
masked chafer beetles, 899
Cyclocephala lurida, 425, 899
ants, 425
Cyclocephala borealis, 899
Cyclocephala hirta, 899
Cyclocephala longula, 899
Cyclocephala pasadenae, 899
masked chafer beetles, 899
Popillia japonica, 425
predators, 425
Solenopsis molesta, 425
turfgrass, 425
Cyclocephala pasadenae,
Cyclocephala borealis, 899
Cyclocephala hirta, 899
Cyclocephala longula, 899
Cyclocephala lurida, 899
masked chafer beetles, 899

Cydia pomonella,
apple, 763
fruit damage, 763
inherited sterility, 763
Cydia trasias,
mating disruption, 7
sex pheromone, 7
tortrix, 7
Cynipidae,
Callirhytis cornigera, 280
density-dependent parasitism, 280
galls, 280
spatial distribution, 280
Cyphocleonus achates,
Agapeta zoegana, 589
biological control, 589
Centaurea maculosa, 589
larval sampling, 589
spotted knapweed, 589

D

Danaus plexippus,
Bacillus thuringiensis, 495
Bt-corn, 495
monarch butterfly, 495
nontarget effects, 495
transgenic, 495
deciduous forests,
beetles, 484
disturbance, 484
forest floor, 484
richness, 484
Shannon diversity index, 484
degradation,
Bacillus thuringiensis, 642
Cry1F, 642
soil, 642
degree-day models,
developmental models, 692
grape berry moth, 692
temperature-dependent growth, 692
degree-days,
development, 23
flight, 23
phenological model, 23
Sesamia nonagrioides, 23
temperature, 23
Dendroctonus pseudotsugae,
bark beetles, 215
disturbance, 215
Douglas-fir beetle, 215
hazard rating model, 215
risk classification model, 215
density, 919, 1047
alatae, 1047
apterae, 1047
development, 1047
egg galleries, 919
forest structure, 919
individual quality, 919
Ips pini, 919
temperature, 919
Toxoptera citricida, 1047
density dependence,
Coleoptera: Silvanidae population regulation, 1061
habitat deterioration, 1061
Oryzaephilus surinamensis, 1061
density-dependent parasitism,
Callirhytis cornigera, 280
Cynipidae, 280
galls, 280
spatial distribution, 280
development, 23, 55, 298, 637, 720, 1047, 1082, 1127
alatae, 1047
aphids, 1127
apterae, 1047
Bemisia tabaci, 55, 720
biological control, 637
biotype B, 720
biotype Q, 720
cantaloupe, 55
Carposina sasakii, 298
corn, 1127
corn aphids, 1127
cotton, 55
degree-days, 23
density, 1047
dry cured-ham, 1082
fecundity, 55, 1127
flight, 23
Lysiphlebus testaceipes, 637
models, 1082
mold mite, 1082
parasitoid, 637
pepper, 720
phenological model, 23
Schizaphis graminum, 637
Sesamia nonagrioides, 23
stage emergence models, 298
survival, 55
temperature, 23
temperatures, 1127
thermal thresholds, 1082
Toxoptera citricida, 1047
Tyrophagus putrescentiae, 1082
developmental models,
degree-day models, 692
grape berry moth, 692
temperature-dependent growth, 692
developmental resistance,
Eucalyptus globulus, 1144
herbivory, 1144
heteroblasty, 1144
phase change, 1144
within-plant variation, 1144
Diabrotica undecimpunctata howardi,
antipredator behavior, 327
predation, 327
sexual differences, 327
trophic cascade, 327
Diaeretiella rapae,
Aphelinus albipodus, 578
Aphelinus asychis, 578
aphid parasitoids, 578
biological control, 578
Russian wheat aphid, 578
diapause, 540, 550, 877, 1090
Adelgidae, 877
Anthonomus grandis, 550
cotton pest management, 550
Endopiza viteana, 540
grape berry moth, 540
horn fly, 1090
overwintering survival, 1090
overwintering survivorship, 550
photoperiod, 877
population dynamics, 1090
temperature, 877, 1090
Tsuga spp. aestivation, 877
Diaprepes abbreviatus,
insect traps, 939
Pachnaeus litus, 939
population monitoring, 939
sample size, 939
sampling, 939
Diapriidae spp.,
Musca domestica, 107
Pachycrepoideus vindemiae, 107
Spalangia spp., 107
Stomoxys calcitrans, 107
Trichomalopsis sp., 107
diel periodicity,
apple, 415
biodiversity, 415
biological control, 415
orchard management, 415
diffusion, 517, 1036
Asian longhorned beetle, 1036
Cerambycidae, 1036
invasive species, 1036
mark-recapture, 1036
simulation model, 517
spatial distribution, 517
Tripleurospermum inodorum, 517
Triticum aestivum, 517
diffusion coefficient,
aerial take-off, 1112
infochemicals, 1112
Spider mites, 1112
walking speed, 1112
diflubenzuron,
Copidosoma floridanum, 794
polyembryony, 794

soybean, 794
Diomorus aiolomorphi,
Aiolomorphus rhopaloides, 1098
bamboo, 1098
gall maker, 1098
phenology, 1098
Diprion pini,
foliage dry matter, 1
hatching, 1
larval survival, 1
Pinus sylvestris, 1
water stress, 1
dispersal, 225, 267, 318, 700, 883
abundance, 267
Acleris gloverana, 883
age-specific survival ratios, 883
aggregation sites, 700
agroecosystems, 225
boundaries, 225
Douglas-fir tussock moth, 883
habitats, 267
highways, 267
mate seeking, 700
mite management, 318
multivariate analysis, 225
Ostrinia nubilalis, 700
Phyllocoptruta oleivora, 318
population dynamics, 883
populations, 267
predators, 225
refuge, 700
Solenopsis invicta, 267
surface airflow, 700
traps, 318
Triticum aestivum, 225
western spruce budworm, 883
distribution, 568, 645
Amblyomma americanum, 568
biogeographical range, 645
habitats, 568
Ixodes scapularis, 568
modeling, 645
quarantine, 645
Solenopsis invicta, 645
ticks, 568
United States, 645
disturbance, 215, 484
bark beetles, 215
beetles, 484
deciduous forests, 484
Dendroctonus pseudotsugae, 215
Douglas-fir beetle, 215
forest floor, 484
hazard rating model, 215
richness, 484
risk classification model, 215
Shannon diversity index, 484
diversity,
ambushers, 1021
hunting spiders, 1021
predatory arthropods, 1021
sampling methods, 1021
web spiders, 1021
diversity indices,
dung beetles, 474
functional groups, 474
habitat change, 474
rank-abundance, 474
size, 474
DNA fingerprinting,
Coptotermes formosanus, 262
genetic structure, 262
dormant spray,
almond, 70
esfenvalerate, 70
Galendromus occidentalis, 70
permethrin, 70
residue, 70
Douglas-fir beetle,
bark beetles, 215
Dendroctonus pseudotsugae, 215
disturbance, 215
hazard rating model, 215
risk classification model, 215
Douglas-fir tussock moth,
Acleris gloverana, 883
age-specific survival ratios, 883
dispersal, 883
population dynamics, 883
western spruce budworm, 883
dry bean,
Epilachna varivestis, 235
life tables, 235
survivorship, 235
dry cured-ham,
development, 1082
models, 1082
mold mite, 1082
thermal thresholds, 1082
Tyrophagus putrescentiae, 1082
Dryas iulia,
body water content, 631
cuticular permeability, 631
Heliconius, 631
surface area, 631
dung beetles,
diversity indices, 474
functional groups, 474
habitat change, 474
rank-abundance, 474
size, 474

E

ectomycorrhiza,
carbon-nutrient balance, 849
gypsy moth, 849
herbivore-plant interactions, 849
Pisolithus tinctorius, 849
Quercus, 849
efficient natural control,
egg parasitism, 37
Geometridae, 37
Lambdina fiscellaria, 37
outbreak collapse, 37
Telenomus, 37
egg galleries,
density, 919
forest structure, 919
individual quality, 919
Ips pini, 919
temperature, 919
egg mass,
beet armyworm, 511
cotton, 511
oviposition, 511
pigweed, 511
Spodoptera exigua, 511
egg parasitism,
efficient natural control, 37
Geometridae, 37
Lambdina fiscellaria, 37
outbreak collapse, 37
Telenomus, 37
eicosanoid,
biocontrol, 449
Coptotermes formosanus, 449
immunity, 449
insect pathology, 449
Serratia marcescens, 449
elm,
feeding injury, 533
maple, 533
nymphal survivorship, 533
oviposition, 533
potato leafhopper, 533
emergence models,
blueberry maggot fly, 925
highbush blueberries, 925
postdiapause development, 925
Rhagoletis mendax, 925
emigration,
Acrolepiopsis assectella, 288
behavior, 288
intercrop, 288
oviposition, 288
Plutella xylostella, 288
Empoasca fabae,
cultural control, 204
intercropping, 204
resource concentration hypothesis, 204
soybean, 204
wheat, 204
Endopiza viteana,
diapause, 540
grape berry moth, 540
Enoclerus sphegeus,
bark beetles, 993
kairomone, 993
Pityogenes carinulatus, 993

Temnochila chlorodia, 993
Thanasimus undatulus, 993
entomopathogenic,
fungi, 196
integrated pest management, 196
nematodes, 196
nontarget organisms, 196
organophosphate pesticides, 196
entomopathogenic nematode,
population dynamics, 254
spatial pattern, 254
environmental risk assessment,
Anticarsia gemmatalis, 1164
beneficial nontarget organisms, 1164
biological control, 1164
biopesticide, 1164
Podisus nigrispinus, 1164
environmental safety,
acetylcholinesterase, 803
bark beetles, 803
carbaryl, 803
Pinaceae, 803
Scolytidae, 803
environmental variables,
Benin, 1135
flight activity, 1135
modeling, 1135
pheromone trapping, 1135
Prostephanus truncatus, 1135
enzyme-linked immunosorbent assay,
bollworm, 431
cotton, 431
Orius, 431
predation, 431
serological assay, 431
Eoreuma loftini,
Allorhogas pyralophagus, 129
functional response, 129
reproduction, 129
Epilachna varivestis, 235, 244
dry bean, 235
geostatistics, 244
life tables, 235
probability distribution, 244
sampling, 244
survivorship, 235
esfenvalerate,
almond, 70
dormant spray, 70
Galendromus occidentalis, 70
permethrin, 70
residue, 70
Eucalyptus globulus, 365, 1144
colored sticky traps, 365
developmental resistance, 1144
herbivory, 1144
heteroblastic host plant, 365
heteroblasty, 1144
phase change, 1144
Psyllid, 365
within-plant variation, 1144
Eurema lisa,
Agraulis vanillae, 1052
Junonia coenia, 1052
phenology, 1052
Phoebis sennae, 1052
Urbanus proteus, 1052
Eurosta solidaginis,
freeze tolerance, 12
over-wintering, 12
supercooling, 12
water balance, 12
Eustenopus villosus,
Asteraceae, 439
biological control, 439
Centaurea solstitialis, 439
Curculionidae, 439
phenological synchrony, 439
extrafloral nectar,
Chrysoperla plorabunda, 595
generalist predator, 595
omnivory, 595

F

fat content,
overwintering, 166
photoperiod, 166
Reticulitermes flavipes, 166
thermoperiod, 166
water content, 166
fatty acids,
Acyrthosiphon pisum, 964
Coccinellidae, 964
Coleomegilla maculata, 964
Hippodamia convergens, 964
tritrophic interactions, 964
fecundity, 55, 1127
aphids, 1127
Bemisia tabaci, 55
cantaloupe, 55
corn, 1127
corn aphids, 1127
cotton, 55
development, 55, 1127
survival, 55
temperatures, 1127
feeding behavior,
herbivory, 172
Japanese beetle, 172
Popillia japonica, 172
thermoregulation, 172
feeding injury,
elm, 533
maple, 533
nymphal survivorship, 533
oviposition, 533
potato leafhopper, 533
female flight propensity,
gypsy moth, 380
hybrids, 380
Lymantria dispar, 380
migration, 380
rhythm, 380
fertilizer,
nitrogen, 501
planting date, 501
population dynamics, 501
field dispersal,
flight characteristics, 1150
flight-mill, 1150
Hyphantria cunea, 1150
mating activity of males, 1150
sex pheromone, 1150
fitness,
genetically modified organism, 972
intra-host competition, 972
mixed infections, 972
nucleopolyhedrovirus, 972
risk assessment, 972
flies,
Rhagoletis cingulata, 82
Rhagoletis fausta, 82
traps, 82
flight, 23, 306
colonization, 306
degree-days, 23
development, 23
impact traps, 306
Leptinotarsa decemlineata, 306
phenological model, 23
plastic-lined trenches, 306
Sesamia nonagrioides, 23
temperature, 23
walking, 306
flight activity,
Benin, 1135
environmental variables, 1135
modeling, 1135
pheromone trapping, 1135
Prostephanus truncatus, 1135
flight behavior,
pest management, 394
tomato yellow leaf curl virus, 394
ultraviolet light, 394
whiteflies, 394
flight characteristics,
field dispersal, 1150
flight-mill, 1150
Hyphantria cunea, 1150
mating activity of males, 1150
sex pheromone, 1150
flight patterns,
insect-plant interactions, 909
Scolytidae, 909
Xyleborinus saxeseni, 909
Xylosandrus crassiusculus, 909
Xylosandrus germanus, 909

flight-mill,
field dispersal, 1150
flight characteristics, 1150
Hyphantria cunea, 1150
mating activity of males, 1150
sex pheromone, 1150
foliage dry matter,
Diprion pini, 1
hatching, 1
larval survival, 1
Pinus sylvestris, 1
water stress, 1
food volatiles,
Attractants, 157
isobutanol, 157
Vespidae, 157
Vespula germanica, 157
yellowjackets, 157
foraging, 123, 673, 832
bone pile, 123
landscape mulch, 832
landscapes, 673
natural enemies, 123
predator, 123
red imported fire ant, 123
refuse pile, 123
resource partitioning, 673
Reticulitermes, 832
Reticulitermes flavipes, 673
Reticulitermes hageni, 673
subterranean termites, 673, 832
foraging behavior, 861, 1103
Hopliinae, 861
insect vision, 861
mimicry, 861
plant-pollinator interactions, 861
predaceous mites, 1103
predator specialization, 1103
prey switching, 1103
Tetranychus urticae, 1103
forest,
ants, 842
Formicidae, 842
habitat, 842
forest floor,
beetles, 484
deciduous forests, 484
disturbance, 484
richness, 484
Shannon diversity index, 484
forest management,
clearcut, 400
insect sampling, 400
selection, 400
shelterwood, 400
vertical distribution of insect fauna, 400
forest structure,
density, 919
egg galleries, 919
individual quality, 919
Ips pini, 919
temperature, 919
Formica,
ant-homopteran interaction, 757
leafhoppers, 757
mutualism, 757
myrmecophile, 757
Formicidae,
ants, 842
forest, 842
habitat, 842
Frankliniella occidentalis,
greenhouse cucumber, 1073
population dynamics, 1073
simulation model, 1073
Franklinothrips orizabensis,
Oligonychus perseae, 892
Persea americana, 892
population dynamics, 892
Scirtothrips perseae, 892
freeze tolerance,
Eurosta solidaginis, 12
over-wintering, 12
supercooling, 12
water balance, 12
frogs,
aggregation, 51
ambush predator, 51
chemical defense, 51
warning coloration, 51
fruit damage,
apple, 763
Cydia pomonella, 763
inherited sterility, 763
functional groups,
diversity indices, 474
dung beetles, 474
habitat change, 474
rank-abundance, 474
size, 474
functional response, 129, 785, 1182
adult size, 1182
Allorhogas pyralophagus, 129
aphid prey, 1182
Aphis gossypii, 785
biological control, 785
coccinellid predator, 1182
Coccinellidae, 785
Eoreuma loftini, 129
larval food supply, 1182
Menochilus sexmaculatus, 1182
population dynamics, 785
reproduction, 129
Scymnus creperus, 785
fungi,
entomopathogenic, 196
integrated pest management, 196
nematodes, 196
nontarget organisms, 196
organophosphate pesticides, 196
fungus,
bacteria, 443
bio-control, 443
Coptotermes formosanus, 443
Serratia marcescens, 443
termite, 443

G

Galendromus occidentalis,
almond, 70
dormant spray, 70
esfenvalerate, 70
permethrin, 70
residue, 70
gall maker,
Aiolomorphus rhopaloides, 1098
bamboo, 1098
Diomorus aiolomorphi, 1098
phenology, 1098
galls,
Callirhytis cornigera, 280
Cynipidae, 280
density-dependent parasitism, 280
spatial distribution, 280
generalist predator,
Chrysoperla plorabunda, 595
extrafloral nectar, 595
omnivory, 595
genetic structure,
Coptotermes formosanus, 262
DNA fingerprinting, 262
genetically modified organism,
fitness, 972
intra-host competition, 972
mixed infections, 972
nucleopolyhedrovirus, 972
risk assessment, 972
geographic information system,
Alphitobius diaperinus, 556
broiler production, 556
lesser mealworm, 556
spatial model, 556
Geometridae,
efficient natural control, 37
egg parasitism, 37
Lambdina fiscellaria, 37
outbreak collapse, 37
Telenomus, 37
geostatistics, 244, 350
Colorado potato beetle, 350
Epilachna varivestis, 244
global positioning systems, 350
insect spatial dynamics, 350
population dynamics, 350
precision agriculture, 350
probability distribution, 244
sampling, 244

global positioning systems,
Colorado potato beetle, 350
geostatistics, 350
insect spatial dynamics, 350
population dynamics, 350
precision agriculture, 350
global warming,
Aedes aegypti, 855
Aedes albopictus, 855
larval development, 855
seasonal environment, 855
survival, 855
Glycine max,
Anticarsia gemmatalis, 76
cultural control, 76
Pseudoplusia includens, 76
Spissistilus festinus, 76
stink bugs, 76
Gracillariidae,
Micrurapteryx salicifoliella, 736
Salix, 736
susceptibility, 736
grape berry moth, 540, 692
degree-day models, 692
developmental models, 692
diapause, 540
Endopiza viteana, 540
temperature-dependent growth, 692
greenhouse cucumber,
Frankliniella occidentalis, 1073
population dynamics, 1073
simulation model, 1073
gypsy moth, 380, 849
carbon-nutrient balance, 849
ectomycorrhiza, 849
female flight propensity, 380
herbivore-plant interactions, 849
hybrids, 380
Lymantria dispar, 380
migration, 380
Pisolithus tinctorius, 849
Quercus, 849
rhythm, 380

H

habitat,
ants, 842
forest, 842
Formicidae, 842
habitat change,
diversity indices, 474
dung beetles, 474
functional groups, 474
rank-abundance, 474
size, 474
habitat deterioration,
Coleoptera: Silvanidae population regulation, 1061
density dependence, 1061
Oryzaephilus surinamensis, 1061
habitat selection,
Aphididae, 371
migratory flight, 371
suction trap, 371
habitats, 267, 568
abundance, 267
Amblyomma americanum, 568
dispersal, 267
distribution, 568
highways, 267
Ixodes scapularis, 568
populations, 267
Solenopsis invicta, 267
ticks, 568
hatching,
Diprion pini, 1
foliage dry matter, 1
larval survival, 1
Pinus sylvestris, 1
water stress, 1
hazard rating model,
bark beetles, 215
Dendroctonus pseudotsugae, 215
disturbance, 215
Douglas-fir beetle, 215
risk classification model, 215
Heliconius,
body water content, 631
cuticular permeability, 631
Dryas iulia, 631
surface area, 631
herbivore-plant interactions,
carbon-nutrient balance, 849
ectomycorrhiza, 849
gypsy moth, 849
Pisolithus tinctorius, 849
Quercus, 849
herbivory, 172, 333, 1144
Asclepias, 333
developmental resistance, 1144
Eucalyptus globulus, 1144
feeding behavior, 172
heteroblasty, 1144
Japanese beetle, 172
phase change, 1144
Popillia japonica, 172
subterranean herbivory, 333
Tetraopes, 333
thermoregulation, 172
within-plant variation, 1144
heteroblastic host plant,
colored sticky traps, 365
Eucalyptus globulus, 365
Psyllid, 365
heteroblasty,
developmental resistance, 1144
Eucalyptus globulus, 1144
herbivory, 1144
phase change, 1144
within-plant variation, 1144
highbush blueberries,
blueberry maggot fly, 925
emergence models, 925
postdiapause development, 925
Rhagoletis mendax, 925
highways,
abundance, 267
dispersal, 267
habitats, 267
populations, 267
Solenopsis invicta, 267
Hippodamia convergens, 811, 964, 1012
Acyrthosiphon pisum, 964
behavior, 1012
biological control, 1012
calorespirometry, 811
Coccinellidae, 964
Coleomegilla maculata, 964
convergent lady beetle, 811
fatty acids, 964
isothermal calorimetry, 811
Myzus nicotianae, 1012
olfactory cues, 1012
prey searching, 1012
respiration, 811
temperature response, 811
tritrophic interactions, 964
Hopliinae,
foraging behavior, 861
insect vision, 861
mimicry, 861
plant-pollinator interactions, 861
horn fly,
diapause, 1090
overwintering survival, 1090
population dynamics, 1090
temperature, 1090
host age,
house fly, 346
parasitization site, 346
parasitoid wasp, 346
pteromalid, 346
Spalangia endius, 346
host damage,
pheromone traps, 776
Pinus taeda, 776
population density, 776
prediction, 776
Rhyacionia frustrana, 776
host plant resistance,
Macrosiphum euphorbiae, 101
Myzus persicae, 101
R genes, 101
resistance management, 101

host range,
Aphantorhaphopsis samarensis, 605
biological control, 605
host specificity, 605
host suitability, 605
host specificity, 605, 742
Aphantorhaphopsis samarensis, 605
Argentine ant, 742
biological control, 605
biological invasion, 742
host range, 605
host suitability, 605
Linepithema humile, 742
Phoridae, 742
Pseudacteon, 742
host stress,
air pollution, 824
Colorado potato beetle, 824
Leptinotarsa decemlineata, 824
ozone, 824
potato, 824
host suitability, 44, 605
Aphantorhaphopsis samarensis, 605
biological control, 605
host range, 605
host specificity, 605
life history, 44
spirea aphid, 44
house fly,
host age, 346
parasitization site, 346
parasitoid wasp, 346
pteromalid, 346
Spalangia endius, 346
hunting spiders,
ambushers, 1021
diversity, 1021
predatory arthropods, 1021
sampling methods, 1021
web spiders, 1021
hybrids,
female flight propensity, 380
gypsy moth, 380
Lymantria dispar, 380
migration, 380
rhythm, 380
Hyphantria cunea,
field dispersal, 1150
flight characteristics, 1150
flight-mill, 1150
mating activity of males, 1150
sex pheromone, 1150

I

immunity,
biocontrol, 449
Coptotermes formosanus, 449
eicosanoid, 449
insect pathology, 449
Serratia marcescens, 449
impact traps,
colonization, 306
flight, 306
Leptinotarsa decemlineata, 306
plastic-lined trenches, 306
walking, 306
individual quality,
density, 919
egg galleries, 919
forest structure, 919
Ips pini, 919
temperature, 919
infochemicals,
aerial take-off, 1112
diffusion coefficient, 1112
spider mites, 1112
walking speed, 1112
inherited sterility,
apple, 763
Cydia pomonella, 763
fruit damage, 763
in-host compatibility,
competition, 1158
Lygus lineolaris, 1158
parasitism, 1158
Peristenus, 1158
tarnished plant bug, 1158
insect pathology,
biocontrol, 449
Coptotermes formosanus, 449
eicosanoid, 449
immunity, 449
Serratia marcescens, 449
insect resistance management,
Bacillus thuringiensis, 409
Ostrinia nubilalis, 409
pollen, 409
transgenic corn, 409
Zea mays, 409
insect sampling,
clearcut, 400
forest management, 400
selection, 400
shelterwood, 400
vertical distribution of insect fauna, 400
insect spatial dynamics,
Colorado potato beetle, 350
geostatistics, 350
global positioning systems, 350
population dynamics, 350
precision agriculture, 350
insect traps,
Diaprepes abbreviatus, 939
Pachnaeus litus, 939
population monitoring, 939
sample size, 939
sampling, 939
insect vision,
foraging behavior, 861
Hopliinae, 861
mimicry, 861
plant-pollinator interactions, 861
insect-plant interactions,
Aleurothrixus aepim, 295
ant-homopteran interactions, 295
Croton floribundus, 295
flight patterns, 909
mutualism, 295
Scolytidae, 909
sooty mold, 295
Xyleborinus saxeseni, 909
Xylosandrus crassiusculus, 909
Xylosandrus germanus, 909
insidious flower bug,
Bacillus thuringiensis, 625
corn, 625
Orius insidiosus, 625
predator, 625
integrated pest management,
entomopathogenic, 196
fungi, 196
nematodes, 196
nontarget organisms, 196
organophosphate pesticides, 196
intercrop,
Acrolepiopsis assectella, 288
behavior, 288
emigration, 288
oviposition, 288
Plutella xylostella, 288
intercropping, 89, 204
Bemisia tabaci, 89
cultural control, 204
Empoasca fabae, 204
polyculture, 89
resource concentration hypothesis, 204
soybean, 204
Trialeurodes vaporariorum, 89
vector management, 89
wheat, 204
intraguild predation,
Cerambycidae, 17
Ips, 17
Ips calligraphus, 17
Monochamus carolinensis, 17
Scolytidae, 17
intra-host competition,
fitness, 972
genetically modified organism, 972
mixed infections, 972
nucleopolyhedrovirus, 972
risk assessment, 972
invading species,
cold hardiness, 181
range limit, 181
southwestern corn borer, 181
winter mortality, 181

invasive pests,
Caribbean region, 339
Maconellicoccus hirsutus, 339
quarantine, 339
sex pheromones, 339
invasive species,
Asian longhorned beetle, 1036
Cerambycidae, 1036
diffusion, 1036
mark-recapture, 1036
Ips,
Cerambycidae, 17
intraguild predation, 17
Ips calligraphus, 17
Monochamus carolinensis, 17
Scolytidae, 17
Ips calligraphus,
Cerambycidae, 17
intraguild predation, 17
Ips, 17
Monochamus carolinensis, 17
Scolytidae, 17
Ips pini,
density, 919
egg galleries, 919
forest structure, 919
individual quality, 919
temperature, 919
isobutanol,
attractants, 157
food volatiles, 157
Vespidae, 157
Vespula germanica, 157
yellowjackets, 157
isothermal calorimetry,
calorespirometry, 811
convergent lady beetle, 811
Hippodamia convergens, 811
respiration, 811
temperature response, 811
Ixodes scapularis,
Amblyomma americanum, 568
distribution, 568
habitats, 568
ticks, 568

J

Japanese beetle,
feeding behavior, 172
herbivory, 172
Popillia japonica, 172
thermoregulation, 172
Junonia coenia,
Agraulis vanillae, 1052
Eurema lisa, 1052
phenology, 1052
Phoebis sennae, 1052
Urbanus proteus, 1052

K

kairomone,
bark beetles, 993
Enoclerus sphegeus, 993
Pityogenes carinulatus, 993
Temnochila chlorodia, 993
Thanasimus undatulus, 993

L

Lacanobia subjuncta,
acetic acid, 656
attractant, 656
Mamestra configurata, 656
trapping, 656
Xestia c-nigrum, 656
lady beetle,
Brassicaceae, 947
Myzus persicae, 947
tritrophic interactions, 947
Lagria hirta,
Coleoptera, 686
Lagriidae, 686
life-history trait correlation, 686
reproduction, 686
Lagriidae,
Coleoptera, 686
Lagria hirta, 686
life-history trait correlation, 686
reproduction, 686
Lambdina fiscellaria,
efficient natural control, 37
egg parasitism, 37
Geometridae, 37
outbreak collapse, 37
Telenomus, 37
landscape mulch,
foraging, 832
Reticulitermes, 832
subterranean termites, 832
landscapes,
foraging, 673
resource partitioning, 673
Reticulitermes flavipes, 673
Reticulitermes hageni, 673
subterranean termites, 673
larval development,
Aedes aegypti, 855
Aedes albopictus, 855
global warming, 855
seasonal environment, 855
survival, 855
larval food supply,
adult size, 1182
aphid prey, 1182
coccinellid predator, 1182
functional response, 1182
Menochilus sexmaculatus, 1182
larval performance,
Chrysomela scripta, 748
Populus, 748
short-rotation forestry, 748
larval sampling,
Agapeta zoegana, 589
biological control, 589
Centaurea maculosa, 589
Cyphocleonus achates, 589
spotted knapweed, 589
larval survival, 1, 728
Bacillus thuringiensis, 728
corn, 728
Diprion pini, 1
foliage dry matter, 1
hatching, 1
Ostrinia nubilalis, 728
Pinus sylvestris, 1
resistance management, 728
water stress, 1
weeds, 728
leafhoppers,
ant–homopteran interaction, 757
Formica, 757
mutualism, 757
myrmecophile, 757
Leptinotarsa decemlineata, 306, 824
air pollution, 824
colonization, 306
Colorado potato beetle, 824
flight, 306
host stress, 824
impact traps, 306
ozone, 824
plastic-lined trenches, 306
potato, 824
walking, 306
Leptura obliterata,
microencapsulation, 837
Pinus ponderosa, 837
plantations, 837
red turpentine beetle, 837
semiochemicals, 837
lesser mealworm,
Alphitobius diaperinus, 556
broiler production, 556
geographic information system, 556
spatial model, 556
life history, 44, 147, 982
biological control, 147, 982
host suitability, 44
Musca domestica, 147, 982
Muscidifurax raptorellus, 982
Muscidifurax zaraptor, 147
spirea aphid, 44
Stomoxys calcitrans, 147, 982
temperature, 147, 982
life table,
bollworm, 112
cotton, 112

ife tables,
dry bean, 235
Epilachna varivestis, 235
survivorship, 235
ife-history trait correlation,
Coleoptera, 686
Lagria hirta, 686
Lagriidae, 686
reproduction, 686
Linepithema humile,
Argentine ant, 742
biological invasion, 742
host specificity, 742
Phoridae, 742
Pseudacteon, 742
Lissorhoptrus oryzophilus,
aquatic barrier trap, 770
aquatic intercept trap, 770
rice integrated pest management, 770
rice water weevil, 770
weevil trapping, 770
Lygus lineolaris,
competition, 1158
in-host compatibility, 1158
parasitism, 1158
Peristenus, 1158
tarnished plant bug, 1158
Lymantria dispar, 380, 1174
Bacillus thuringiensis, 1174
Compsilura, 1174
female flight propensity, 380
gypsy moth, 380
hybrids, 380
migration, 380
rhythm, 380
sublethal effects, 1174
Lysiphlebus testaceipes,
biological control, 637
development, 637
parasitoid, 637
Schizaphis graminum, 637

M

Maconellicoccus hirsutus,
Caribbean region, 339
invasive pests, 339
quarantine, 339
sex pheromones, 339
Macrosiphum euphorbiae,
host plant resistance, 101
Myzus persicae, 101
R genes, 101
resistance management, 101
Mamestra configurata,
acetic acid, 656
attractant, 656
Lacanobia subjuncta, 656
trapping, 656
Xestia c-nigrum, 656
maple,
elm, 533
feeding injury, 533
nymphal survivorship, 533
oviposition, 533
potato leafhopper, 533
marking,
arthropods, 711
movement, 711
predator, 711
rubidium, 711
mark-recapture,
Asian longhorned beetle, 1036
Cerambycidae, 1036
diffusion, 1036
invasive species, 1036
masked chafer beetles,
Cyclocephala borealis, 899
Cyclocephala hirta, 899
Cyclocephala longula, 899
Cyclocephala lurida, 899
Cyclocephala pasadenae, 899
mate seeking,
aggregation sites, 700
dispersal, 700
Ostrinia nubilalis, 700
refuge, 700
surface airflow, 700
mating activity of males,
field dispersal, 1150
flight characteristics, 1150
flight-mill, 1150
Hyphantria cunea, 1150
sex pheromone, 1150
mating disruption,
Cydia trasias, 7
sex pheromone, 7
tortrix, 7
mating success,
multiple mating, 31
Phthorimaea operculella, 31
Menochilus sexmaculatus,
adult size, 1182
aphid prey, 1182
coccinellid predator, 1182
functional response, 1182
larval food supply, 1182
microencapsulation,
Leptura obliterata, 837
Pinus ponderosa, 837
plantations, 837
red turpentine beetle, 837
semiochemicals, 837
Microtrombidiidae,
Brachycera, 903
Parasitization, 903
Tachinidae, 903
Therevidae, 903
Micrurapteryx salicifoliella,
Gracillariidae, 736
Salix, 736
susceptibility, 736
migration, 189, 380
black cutworm moth, 189
calling, 189
female flight propensity, 380
gypsy moth, 380
hybrids, 380
Lymantria dispar, 380
periodicity, 189
reproductive diapause, 189
rhythm, 380
sex pheromone, 189
migratory flight,
Aphididae, 371
habitat selection, 371
suction trap, 371
mimicry,
foraging behavior, 861
Hopliinae, 861
insect vision, 861
plant-pollinator interactions, 861
mite management,
dispersal, 318
Phyllocoptruta oleivora, 318
traps, 318
mixed infections,
fitness, 972
genetically modified organism, 972
intra-host competition, 972
nucleopolyhedrovirus, 972
risk assessment, 972
modeling, 645, 1135
Benin, 1135
biogeographical range, 645
distribution, 645
environmental variables, 1135
flight activity, 1135
pheromone trapping, 1135
Prostephanus truncatus, 1135
quarantine, 645
Solenopsis invicta, 645
United States, 645
models,
development, 1082
dry cured-ham, 1082
mold mite, 1082
thermal thresholds, 1082
Tyrophagus putrescentiae, 1082
mold mite,
development, 1082
dry cured-ham, 1082
models, 1082
thermal thresholds, 1082
Tyrophagus putrescentiae, 1082
mole crickets,
behavior, 466
oviposition, 466
Scapteriscus borellii, 466
soil moisture, 466

monarch butterfly,
Bacillus thuringiensis, 495
Bt-corn, 495
Danaus plexippus, 495
nontarget effects, 495
transgenic, 495
Monochamus carolinensis,
Cerambycidae, 17
intraguild predation, 17
Ips, 17
Ips calligraphus, 17
Scolytidae, 17
mortality,
Choristoneura occidentalis, 136
multicapsid nucleopolyhedrovirus, 136
sublethal effect, 136
mosquito,
Aedes sollicitans, 388
control, 388
starch, 388
xanthan gum, 388
movement,
arthropods, 711
marking, 711
predator, 711
rubidium, 711
multicapsid nucleopolyhedrovirus,
Choristoneura occidentalis, 136
mortality, 136
nucleopolyhedrovirus, 136
sublethal effect, 136
multiple mating,
mating success, 31
Phthorimaea operculella, 31
multiple regression stepwise selection procedure analysis,
accumulated air mean subfreezing temperature, 562
bean leaf beetle, 562
Cerotoma trifurcata, 562
population dynamics, 562
multivariate analysis,
agroecosystems, 225
boundaries, 225
dispersal, 225
predators, 225
Triticum aesticum, 225
Musca domestica, 107, 147, 982
biological control, 147, 982
Diapriidae spp., 107
life history, 147, 982
Muscidifurax raptorellus, 982
Muscidifurax zaraptor, 147
Pachycrepoideus vindemiae, 107
Spalangia spp., 107
Stomoxys calcitrans, 107, 147, 982
temperature, 147, 982
Trichomalopsis sp., 107
Muscidifurax raptorellus,
biological control, 982
life history, 982
Musca domestica, 982
Stomoxys calcitrans, 982
temperature, 982
Muscidifurax zaraptor,
biological control, 147
life history, 147
Musca domestica, 147
Stomoxys calcitrans, 147
temperature, 147
musk thistle,
biological control, 612
Carduus nutans, 612
red imported fire ant, 612
Rhinocyllus conicus, 612
Solenopsis invicta, 612
mutualism, 295, 757
Aleurothrixus aepim, 295
ant–homopteran interaction, 757
ant-homopteran interactions, 295
Croton floribundus, 295
Formica, 757
insect–plant interactions, 295
leafhoppers, 757
myrmecophile, 757
sooty mold, 295
mycotoxins,
aflatoxins, 817
protein metabolism, 817
Spodoptera littoralis, 817
myrmecophile,
ant–homopteran interaction, 757
Formica, 757
leafhoppers, 757
mutualism, 757
Myzus nicotianae,
behavior, 1012
biological control, 1012
Hippodamia convergens, 1012
olfactory cues, 1012
prey searching, 1012
Myzus persicae, 101, 947
Brassicaceae, 947
host plant resistance, 101
lady beetle, 947
Macrosiphum euphorbiae, 101
R genes, 101
resistance management, 101
tritrophic interactions, 947

N

natural enemies,
bone pile, 123
foraging, 123
predator, 123
red imported fire ant, 123
refuse pile, 123
nematodes,
entomopathogenic, 196
fungi, 196
integrated pest management, 196
nontarget organisms, 196
organophosphate pesticides, 196
Nezara viridula,
Anoplolepis longipes, 1028
biological control, 1028
community structure, 1028
Pheidole megacephala, 1028
nitrogen,
fertilizer, 501
planting date, 501
population dynamics, 501
nontarget effects,
Bacillus thuringiensis, 495
Bt-corn, 495
Danaus plexippus, 495
monarch butterfly, 495
transgenic, 495
nontarget impact,
biological control of weeds, 953
Carthamus tinctorius, 953
Centaurea solstitialis, 953
choice tests, 953
realized host range, 953
nontarget organisms,
entomopathogenic, 196
fungi, 196
integrated pest management, 196
nematodes, 196
organophosphate pesticides, 196
nucleopolyhedrovirus, 136, 972
Choristoneura occidentalis, 136
fitness, 972
genetically modified organism, 972
intra-host competition, 972
mixed infections, 972
mortality, 136
multicapsid nucleopolyhedrovirus, 136
risk assessment, 972
sublethal effect, 136
nymphal survivorship,
elm, 533
feeding injury, 533
maple, 533

oviposition, 533
potato leafhopper, 533

O

O_3,
Aspen-FACE, 1119
Betula papyrifera, 1119
CO_2, 1119
Orgyia leucostigma whitemarked tussock moth, 1119
olfactory cues,
behavior, 1012
biological control, 1012
Hippodamia convergens, 1012
Myzus nicotianae, 1012
prey searching, 1012
Oligonychus perseae,
Franklinothrips orizabensis, 892
Persea americana, 892
population dynamics, 892
Scirtothrips perseae, 892
omnivory,
Chrysoperla plorabunda, 595
extrafloral nectar, 595
generalist predator, 595
orchard management,
apple, 415
biodiversity, 415
biological control, 415
diel periodicity, 415
organophosphate pesticides,
entomopathogenic, 196
fungi, 196
integrated pest management, 196
nematodes, 196
nontarget organisms, 196
Orgyia leucostigma whitemarked tussock moth,
Aspen-FACE, 1119
Betula papyrifera, 1119
CO_2, 1119
O_3, 1119
Orius,
bollworm, 431
cotton, 431
enzyme-linked immunosorbent assay, 431
predation, 431
serological assay, 431
Orius insidiosus,
Bacillus thuringiensis, 625
corn, 625
insidious flower bug, 625
predator, 625
Oryzaephilus surinamensis, 617, 1061
biological control, 617
Coleoptera: Silvanidae population regulation, 1061
density dependence, 1061
habitat deterioration, 1061
Plodia interpunctella, 617
Rhyzopertha dominica, 617
stored products, 617
Tribolium castaneum, 617
Ostrinia nubilalis, 274, 409, 700, 728
aggregation sites, 700
Bacillus thuringiensis, 409, 728
corn, 728
dispersal, 700
insect resistance management, 409
larval survival, 728
mate seeking, 700
oviposition, 274
pollen, 409
refuge, 700
resistance management, 728
surface airflow, 700
sweet corn, 274
transgenic corn, 409
weeds, 728
Zea mays, 274, 409
outbreak collapse,
efficient natural control, 37
egg parasitism, 37
Geometridae, 37
Lambdina fiscellaria, 37
Telenomus, 37
overwintering, 64, 166
boll weevil, 64
cultural management, 64
Eurosta solidaginis, 12
fat content, 166
freeze tolerance, 12
pheromone synergist, 64
photoperiod, 166
Reticulitermes flavipes, 166
supercooling, 12
thermoperiod, 166
trapping, 64
water balance, 12
water content, 166
overwintering survival,
diapause, 1090
horn fly, 1090
population dynamics, 1090
temperature, 1090
overwintering survivorship,
Anthonomus grandis, 550
cotton pest management, 550
diapause, 550
oviposition, 274, 288, 466, 511, 533
Acrolepiopsis assectella, 288
beet armyworm, 511
behavior, 288, 466
cotton, 511
egg mass, 511
elm, 533
emigration, 288
feeding injury, 533
intercrop, 288
maple, 533
mole crickets, 466
nymphal survivorship, 533
Ostrinia nubilalis, 274
pigweed, 511
Plutella xylostella, 288
potato leafhopper, 533
Scapteriscus borellii, 466
soil moisture, 466
Spodoptera exigua, 511
sweet corn, 274
Zea mays, 274
ozone,
air pollution, 824
Colorado potato beetle, 824
host stress, 824
Leptinotarsa decemlineata, 824
potato, 824

P

Pachnaeus litus,
Diaprepes abbreviatus, 939
insect traps, 939
population monitoring, 939
sample size, 939
sampling, 939
Pachycrepoideus vindemiae,
Diapriidae spp., 107
Musca domestica, 107
Spalangia spp., 107
Stomoxys calcitrans, 107
Trichomalopsis sp., 107
parasitism,
competition, 1158
in-host compatibility, 1158
Lygus lineolaris, 1158
Peristenus, 1158
tarnished plant bug, 1158
parasitization,
Brachycera, 903
Microtrombidiidae, 903
Tachinidae, 903
Therevidae, 903
parasitization site,
host age, 346
house fly, 346
parasitoid wasp, 346
pteromalid, 346
Spalangia endius, 346
parasitoid, 637, 869
biological control, 637
Cotesia flavipes, 869
development, 637
Lysiphlebus testaceipes, 637
Schizaphis graminum, 637
spatial interpolation, 869
spatial-temporal population dynamics, 869

validation, 869
parasitoid wasp,
host age, 346
house fly, 346
parasitization site, 346
pteromalid, 346
Spalangia endius, 346
pepper,
Bemisia tabaci, 720
biotype B, 720
biotype Q, 720
development, 720
periodicity,
black cutworm moth, 189
calling, 189
migration, 189
reproductive diapause, 189
sex pheromone, 189
Peristenus,
competition, 1158
in-host compatibility, 1158
Lygus lineolaris, 1158
parasitism, 1158
tarnished plant bug, 1158
permethrin,
almond, 70
dormant spray, 70
esfenvalerate, 70
Galendromus occidentalis, 70
residue, 70
Persea americana,
Franklinothrips orizabensis, 892
Oligonychus perseae, 892
population dynamics, 892
Scirtothrips perseae, 892
pest management,
flight behavior, 394
tomato yellow leaf curl virus, 394
ultraviolet light, 394
whiteflies, 394
phase change,
developmental resistance, 1144
Eucalyptus globulus, 1144
herbivory, 1144
heteroblasty, 1144
within-plant variation, 1144
Pheidole megacephala,
Anoplolepis longipes, 1028
biological control, 1028
community structure, 1028
Nezara viridula, 1028
phenological model,
degree-days, 23
development, 23
flight, 23
Sesamia nonagrioides, 23
temperature, 23
phenological synchrony,
Asteraceae, 439
biological control, 439
Centaurea solstitialis, 439
Curculionidae, 439
Eustenopus villosus, 439
phenology, 1052, 1098
Agraulis vanillae, 1052
Aiolomorphus rhopaloides, 1098
bamboo, 1098
Diomorus aiolomorphi, 1098
Eurema lisa, 1052
gall maker, 1098
Junonia coenia, 1052
Phoebis sennae, 1052
Urbanus proteus, 1052
phenylacetaldehyde,
alfalfa looper, 667
attractant, 667
Autographa californica, 667
trap, 667
pheromone quantity,
component ratio, 1006
Rhyacionia frustrana, 1006
sex pheromone, 1006
synthetic bait, 1006
pheromone synergist,
boll weevil, 64
cultural management, 64
overwintering, 64
trapping, 64
pheromone trap catch,
adult longevity, 999
pupal weight, 999
Rhyacionia frustrana, 999
pheromone trapping,
Benin, 1135
environmental variables, 1135
flight activity, 1135
modeling, 1135
Prostephanus truncatus, 1135
pheromone traps,
host damage, 776
Pinus taeda, 776
population density, 776
prediction, 776
Rhyacionia frustrana, 776
Phoebis sennae,
Agraulis vanillae, 1052
Eurema lisa, 1052
Junonia coenia, 1052
phenology, 1052
Urbanus proteus, 1052
Phoridae,
Argentine ant, 742
biological invasion, 742
host specificity, 742
Linepithema humile, 742
Pseudacteon, 742
photoperiod, 166, 877
Adelgidae, 877
diapause, 877
fat content, 166
overwintering, 166
Reticulitermes flavipes, 166
temperature, 877
thermoperiod, 166
Tsuga spp. aestivation, 877
water content, 166
Phthorimaea operculella,
mating success, 31
multiple mating, 31
Phyllocoptruta oleivora,
dispersal, 318
mite management, 318
traps, 318
pigweed,
beet armyworm, 511
cotton, 511
egg mass, 511
oviposition, 511
Spodoptera exigua, 511
Pinaceae,
acetylcholinesterase, 803
bark beetles, 803
carbaryl, 803
environmental safety, 803
Scolytidae, 803
Pinus ponderosa,
Leptura obliterata, 837
microencapsulation, 837
plantations, 837
red turpentine beetle, 837
semiochemicals, 837
Pinus sylvestris,
Diprion pini, 1
foliage dry matter, 1
hatching, 1
larval survival, 1
water stress, 1
Pinus taeda,
host damage, 776
pheromone traps, 776
population density, 776
prediction, 776
Rhyacionia frustrana, 776
Pisolithus tinctorius,
carbon-nutrient balance, 849
ectomycorrhiza, 849
gypsy moth, 849
herbivore-plant interactions, 849
Quercus, 849
Pityogenes carinulatus,
bark beetles, 993
Enoclerus sphegeus, 993
kairomone, 993
Temnochila chlorodia, 993
Thanasimus undatulus, 993
plant resistance,
alternating current electrical penetration graph system, 1041
corn leaf aphid, 1041
plant water stress, 1041
probing behavior, 1041

Russian wheat aphid, 1041
plant water stress,
alternating current electrical penetration graph system, 1041
corn leaf aphid, 1041
plant resistance, 1041
probing behavior, 1041
Russian wheat aphid, 1041
plantations,
Leptura obliterata, 837
microencapsulation, 837
Pinus ponderosa, 837
red turpentine beetle, 837
semiochemicals, 837
planting date,
fertilizer, 501
nitrogen, 501
population dynamics, 501
plant–insect interactions,
Bacillus thuringiensis tenebrionis, 524
Colorado potato beetle, 524
potato, 524
potato aphid, 524
transgenic classical resistance, 524
plant-pollinator interactions,
foraging behavior, 861
Hopliinae, 861
insect vision, 861
mimicry, 861
plastic-lined trenches,
colonization, 306
flight, 306
impact traps, 306
Leptinotarsa decemlineata, 306
walking, 306
Plodia interpunctella,
biological control, 617
Oryzaephilus surinamensis, 617
Rhyzopertha dominica, 617
stored products, 617
Tribolium castaneum, 617
Plutella xylostella,
Acrolepiopsis assectella, 288
behavior, 288
emigration, 288
intercrop, 288
oviposition, 288
Podisus nigrispinus,
Anticarsia gemmatalis, 1164
beneficial nontarget organisms, 1164
biological control, 1164
biopesticide, 1164
environmental risk assessment, 1164
pollen,
Bacillus thuringiensis, 409
insect resistance management, 409
Ostrinia nubilalis, 409
transgenic corn, 409
Zea mays, 409
polyculture,
Bemisia tabaci, 89
intercropping, 89
Trialeurodes vaporariorum, 89
vector management, 89
polyembryony,
Copidosoma floridanum, 794
diflubenzuron, 794
soybean, 794
Popillia japonica, 172, 425
ants, 425
Cyclocephala lurida, 425
feeding behavior, 172
herbivory, 172
Japanese beetle, 172
predators, 425
Solenopsis molesta, 425
thermoregulation, 172
turfgrass, 425
population density,
host damage, 776
pheromone traps, 776
Pinus taeda, 776
prediction, 776
Rhyacionia frustrana, 776
population dynamics, 254, 350, 501, 562, 785, 883, 892, 1073, 1090
accumulated air mean subfreezing temperature, 562
Acleris gloverana, 883
age-specific survival ratios, 883
Aphis gossypii, 785
bean leaf beetle, 562
biological control, 785
Cerotoma trifurcata, 562
Coccinellidae, 785
Colorado potato beetle, 350
diapause, 1090
dispersal, 883
Douglas-fir tussock moth, 883
entomopathogenic nematode, 254
fertilizer, 501
Frankliniella occidentalis, 1073
Franklinothrips orizabensis, 892
functional response, 785
geostatistics, 350
global positioning systems, 350
greenhouse cucumber, 1073
horn fly, 1090
insect spatial dynamics, 350
multiple regression stepwise selection procedure analysis, 562
nitrogen, 501
Oligonychus perseae, 892
overwintering survival, 1090
Persea americana, 892
planting date, 501
precision agriculture, 350
Scirtothrips perseae, 892
Scymnus creperus, 785
simulation model, 1073
spatial pattern, 254
temperature, 1090
western spruce budworm, 883
population monitoring, 663, 939
Diaprepes abbreviatus, 939
insect traps, 939
Pachnaeus litus, 939
sample size, 939
sampling, 939
sex pheromone, 663
sticky traps, 663
Synanthedon tipuliformis, 663
populations,
abundance, 267
dispersal, 267
habitats, 267
highways, 267
Solenopsis invicta, 267
Populus,
Chrysomela scripta, 748
larval performance, 748
short-rotation forestry, 748
postdiapause development,
blueberry maggot fly, 925
emergence models, 925
highbush blueberries, 925
Rhagoletis mendax, 925
potato, 524, 824
air pollution, 824
Bacillus thuringiensis tenebrionis, 524
Colorado potato beetle, 524, 824
host stress, 824
Leptinotarsa decemlineata, 824
ozone, 824
plant–insect interactions, 524
potato aphid, 524
transgenic classical resistance, 524
potato aphid,
Bacillus thuringiensis tenebrionis, 524
Colorado potato beetle, 524
plant–insect interactions, 524
potato, 524
transgenic classical resistance, 524
potato leafhopper,
elm, 533
feeding injury, 533
maple, 533
nymphal survivorship, 533
oviposition, 533

precision agriculture,
Colorado potato beetle, 350
geostatistics, 350
global positioning systems, 350
insect spatial dynamics, 350
population dynamics, 350
predaceous mites,
foraging behavior, 1103
predator specialization, 1103
prey switching, 1103
Tetranychus urticae, 1103
predation, 327, 431
antipredator behavior, 327
bollworm, 431
cotton, 431
Diabrotica undecimpunctata howardi, 327
enzyme-linked immunosorbent assay, 431
Orius, 431
serological assay, 431
sexual differences, 327
trophic cascade, 327
predator, 123, 625, 711
arthropods, 711
Bacillus thuringiensis, 625
bone pile, 123
corn, 625
foraging, 123
insidious flower bug, 625
marking, 711
movement, 711
natural enemies, 123
Orius insidiosus, 625
red imported fire ant, 123
refuse pile, 123
rubidium, 711
predator specialization,
foraging behavior, 1103
predaceous mites, 1103
prey switching, 1103
Tetranychus urticae, 1103
predators, 225, 425
agroecosystems, 225
ants, 425
boundaries, 225
Cyclocephala lurida, 425
dispersal, 225
multivariate analysis, 225
Popillia japonica, 425
Solenopsis molesta, 425
Triticum aestivum, 225
turfgrass, 425
predatory arthropods,
ambushers, 1021
diversity, 1021
hunting spiders, 1021
sampling methods, 1021
web spiders, 1021
prediction,
host damage, 776
pheromone traps, 776
Pinus taeda, 776
population density, 776
Rhyacionia frustrana, 776
prey searching,
behavior, 1012
biological control, 1012
Hippodamia convergens, 1012
Myzus nicotianae, 1012
olfactory cues, 1012
prey switching,
foraging behavior, 1103
predaceous mites, 1103
predator specialization, 1103
Tetranychus urticae, 1103
probability distribution,
Epilachna varivestis, 244
geostatistics, 244
sampling, 244
probing behavior,
alternating current electrical penetration graph system, 1041
corn leaf aphid, 1041
plant resistance, 1041
plant water stress, 1041
Russian wheat aphid, 1041
Prostephanus truncatus,
Benin, 1135
environmental variables, 1135
flight activity, 1135
modeling, 1135
pheromone trapping, 1135
protein metabolism,
aflatoxins, 817
mycotoxins, 817
Spodoptera littoralis, 817
Pseudacteon,
Argentine ant, 742
biological invasion, 742
host specificity, 742
Linepithema humile, 742
Phoridae, 742
Pseudoplusia includens,
Anticarsia gemmatalis, 76
cultural control, 76
Glycine max, 76
Spissistilus festinus, 76
stink bugs, 76
Psyllid,
colored sticky traps, 365
Eucalyptus globulus, 365
heteroblastic host plant, 365
pteromalid,
host age, 346
house fly, 346
parasitization site, 346
parasitoid wasp, 346
Spalangia endius, 346
pupal weight,
adult longevity, 999
pheromone trap catch, 999
Rhyacionia frustrana, 999

Q

quarantine, 339, 645
biogeographical range, 645
Caribbean region, 339
distribution, 645
invasive pests, 339
Maconellicoccus hirsutus, 339
modeling, 645
sex pheromones, 339
Solenopsis invicta, 645
United States, 645
Quercus,
carbon-nutrient balance, 849
ectomycorrhiza, 849
gypsy moth, 849
herbivore-plant interactions, 849
Pisolithus tinctorius, 849

R

R genes,
host plant resistance, 101
Macrosiphum euphorbiae, 101
Myzus persicae, 101
resistance management, 101
Reticulitermes flavipes,
Coptotermes formosanus, 545
termite tunneling, 545
tunneling geometry, 545
range limit,
cold hardiness, 181
invading species, 181
southwestern corn borer, 181
winter mortality, 181
rank-abundance,
diversity indices, 474
dung beetles, 474
functional groups, 474
habitat change, 474
size, 474
realized host range,
biological control of weeds, 953
Carthamus tinctorius, 953
Centaurea solstitialis, 953
choice tests, 953
nontarget impact, 953
red imported fire ant, 123, 612
biological control, 612
bone pile, 123
Carduus nutans, 612
foraging, 123
musk thistle, 612
natural enemies, 123
predator, 123
refuse pile, 123
Rhinocyllus conicus, 612

Solenopsis invicta, 612
red turpentine beetle,
Leptura obliterata, 837
microencapsulation, 837
Pinus ponderosa, 837
plantations, 837
semiochemicals, 837
refuge,
aggregation sites, 700
dispersal, 700
mate seeking, 700
Ostrinia nubilalis, 700
surface airflow, 700
refuse pile,
bone pile, 123
foraging, 123
natural enemies, 123
predator, 123
red imported fire ant, 123
reproduction, 129, 686
Allorhogas pyralophagus, 129
Coleoptera, 686
Eoreuma loftini, 129
functional response, 129
Lagria hirta, 686
Lagriidae, 686
life-history trait correlation, 686
reproductive diapause,
black cutworm moth, 189
calling, 189
migration, 189
periodicity, 189
sex pheromone, 189
residue,
almond, 70
dormant spray, 70
esfenvalerate, 70
Galendromus occidentalis, 70
permethrin, 70
resistance management, 101, 728
Bacillus thuringiensis, 728
corn, 728
host plant resistance, 101
larval survival, 728
Macrosiphum euphorbiae, 101
Myzus persicae, 101
Ostrinia nubilalis, 728
R genes, 101
weeds, 728
resource concentration hypothesis,
cultural control, 204
Empoasca fabae, 204
intercropping, 204
soybean, 204
wheat, 204
resource partitioning,
foraging, 673
landscapes, 673
Reticulitermes flavipes, 673
Reticulitermes hageni, 673
subterranean termites, 673
respiration,
calorespirometry, 811
convergent lady beetle, 811
Hippodamia convergens, 811
isothermal calorimetry, 811
temperature response, 811
Reticulitermes,
foraging, 832
landscape mulch, 832
subterranean termites, 832
Reticulitermes flavipes, 166, 457, 673
chemical communication, 457
Coptotermes formosanus, 457
fat content, 166
foraging, 673
landscapes, 673
overwintering, 166
photoperiod, 166
resource partitioning, 673
Reticulitermes hageni, 673
subterranean termites, 673
thermoperiod, 166
trail pheromone, 457
water content, 166
Reticulitermes hageni,
foraging, 673
landscapes, 673
resource partitioning, 673
Reticulitermes flavipes, 673
subterranean termites, 673
Rhagoletis cingulata,
flies, 82
Rhagoletis fausta, 82
traps, 82
Rhagoletis fausta,
flies, 82
Rhagoletis cingulata, 82
traps, 82
Rhagoletis mendax,
blueberry maggot fly, 925
emergence models, 925
highbush blueberries, 925
postdiapause development, 925
Rhinocyllus conicus,
biological control, 612
Carduus nutans, 612
musk thistle, 612
red imported fire ant, 612
Solenopsis invicta, 612
Rhyacionia frustrana, 776, 999, 1006
adult longevity, 999
component ratio, 1006
host damage, 776
pheromone quantity, 1006
pheromone trap catch, 999
pheromone traps, 776
Pinus taeda, 776
population density, 776
prediction, 776
pupal weight, 999
sex pheromone, 1006
synthetic bait, 1006
rhythm,
female flight propensity, 380
gypsy moth, 380
hybrids, 380
Lymantria dispar, 380
migration, 380
Rhyzopertha dominica,
biological control, 617
Oryzaephilus surinamensis, 617
Plodia interpunctella, 617
stored products, 617
Tribolium castaneum, 617
rice integrated pest management,
aquatic barrier trap, 770
aquatic intercept trap, 770
Lissorhoptrus oryzophilus, 770
rice water weevil, 770
weevil trapping, 770
rice water weevil,
aquatic barrier trap, 770
aquatic intercept trap, 770
Lissorhoptrus oryzophilus, 770
rice integrated pest management, 770
weevil trapping, 770
richness,
beetles, 484
deciduous forests, 484
disturbance, 484
forest floor, 484
Shannon diversity index, 484
risk assessment,
fitness, 972
genetically modified organism, 972
intra-host competition, 972
mixed infections, 972
nucleopolyhedrovirus, 972
risk classification model,
bark beetles, 215
Dendroctonus pseudotsugae, 215
disturbance, 215
Douglas-fir beetle, 215
hazard rating model, 215
rubidium,
arthropods, 711
marking, 711
movement, 711
predator, 711
Russian wheat aphid, 578, 1041
alternating current electrical penetration graph system, 1041
Aphelinus albipodus, 578
Aphelinus asychis, 578
aphid parasitoids, 578
biological control, 578

corn leaf aphid, 1041
Diaeretiella rapae, 578
plant resistance, 1041
plant water stress, 1041
probing behavior, 1041

S

Salix,
Gracillariidae, 736
Micrurapteryx salicifoliella, 736
susceptibility, 736
sample size,
Diaprepes abbreviatus, 939
insect traps, 939
Pachnaeus litus, 939
population monitoring, 939
sampling, 939
sampling, 244, 939
Diaprepes abbreviatus, 939
Epilachna varivestis, 244
geostatistics, 244
insect traps, 939
Pachnaeus litus, 939
population monitoring, 939
probability distribution, 244
sample size, 939
sampling methods,
ambushers, 1021
diversity, 1021
hunting spiders, 1021
predatory arthropods, 1021
web spiders, 1021
Scapteriscus borellii,
behavior, 466
mole crickets, 466
oviposition, 466
soil moisture, 466
scarring on avocado fruit,
avocado, 932
Scirtothrips perseae, 932
thresholds, 932
thrips numbers, 932
Schizaphis graminum,
biological control, 637
development, 637
Lysiphlebus testaceipes, 637
parasitoid, 637
Scirtothrips perseae, 892, 932
avocado, 932
Franklinothrips orizabensis, 892
Oligonychus perseae, 892
Persea americana, 892
population dynamics, 892
scarring on avocado fruit, 932
thresholds, 932
thrips numbers, 932
Scolytidae, 17, 803, 909
acetylcholinesterase, 803
bark beetles, 803
carbaryl, 803
Cerambycidae, 17
environmental safety, 803
flight patterns, 909
insect-plant interactions, 909
intraguild predation, 17
Ips, 17
Ips calligraphus, 17
Monochamus carolinensis, 17
Pinaceae, 803
Xyleborinus saxeseni, 909
Xylosandrus crassiusculus, 909
Xylosandrus germanus, 909
Scymnus creperus,
Aphis gosspii, 785
biological control, 785
Coccinellidae, 785
functional response, 785
population dynamics, 785
seasonal environment,
Aedes aegypti, 855
Aedes albopictus, 855
global warming, 855
larval development, 855
survival, 855
selection,
clearcut, 400
forest management, 400
insect sampling, 400
shelterwood, 400
vertical distribution of insect fauna, 400
semiochemicals,
Leptura obliterata, 837
microencapsulation, 837
Pinus ponderosa, 837
plantations, 837
red turpentine beetle, 837
serological assay,
bollworm, 431
cotton, 431
enzyme-linked immunosorbent assay, 431
Orius, 431
predation, 431
Serratia marcescens, 443, 449
bacteria, 443
bio-control, 443
biocontrol, 449
Coptotermes formosanus, 443, 449
eicosanoid, 449
fungus, 443
immunity, 449
insect pathology, 449
termite, 443
Sesamia nonagrioides,
degree-days, 23
development, 23
flight, 23
phenological model, 23
temperature, 23
sex pheromone, 7, 189, 663, 1006, 1150
black cutworm moth, 189
calling, 189
component ratio, 1006
Cydia trasias, 7
field dispersal, 1150
flight characteristics, 1150
flight-mill, 1150
Hyphantria cunea, 1150
mating activity of males, 1150
mating disruption, 7
migration, 189
periodicity, 189
pheromone quantity, 1006
population monitoring, 663
reproductive diapause, 189
Rhyacionia frustrana, 1006
sticky traps, 663
Synanthedon tipuliformis, 663
synthetic bait, 1006
tortrix, 7
sex pheromones,
Caribbean region, 339
invasive pests, 339
Maconellicoccus hirsutus, 339
quarantine, 339
sexual differences,
antipredator behavior, 327
Diabrotica undecimpunctata howardi, 327
predation, 327
trophic cascade, 327
Shannon diversity index,
beetles, 484
deciduous forests, 484
disturbance, 484
forest floor, 484
richness, 484
shelterwood,
clearcut, 400
forest management, 400
insect sampling, 400
selection, 400
vertical distribution of insect fauna, 400
short-rotation forestry,
Chrysomela scripta, 748
larval performance, 748
Populus, 748
simulation model, 517, 1073
diffusion, 517
Frankliniella occidentalis, 1073
greenhouse cucumber, 1073
population dynamics, 1073
spatial distribution, 517
Tripleurospermum inodorum, 517
Triticum aestivum, 517
size,
diversity indices, 474
dung beetles, 474

functional groups, 474
habitat change, 474
rank-abundance, 474
soil,
Bacillus thuringiensis, 642
Cry1F, 642
degradation, 642
soil moisture,
behavior, 466
mole crickets, 466
oviposition, 466
Scapteriscus borellii, 466
Solenopsis invicta, 267, 612, 645, 1017
abundance, 267
arthropods, 1017
biogeographical range, 645
biological control, 612
Carduus nutans, 612
cover crops, 1017
dispersal, 267
distribution, 645
habitats, 267
highways, 267
modeling, 645
musk thistle, 612
populations, 267
quarantine, 645
red imported fire ant, 612
Rhinocyllus conicus, 612
sugarcane, 1017
United States, 645
Solenopsis molesta,
ants, 425
Cyclocephala lurida, 425
Popillia japonica, 425
predators, 425
turfgrass, 425
sooty mold,
Aleurothrixus aepim, 295
ant-homopteran interactions, 295
Croton floribundus, 295
insect–plant interactions, 295
mutualism, 295
southwestern corn borer,
cold hardiness, 181
invading species, 181
range limit, 181
winter mortality, 181
soybean, 204, 794
Copidosoma floridanum, 794
cultural control, 204
diflubenzuron, 794
Empoasca fabae, 204
intercropping, 204
polyembryony, 794
resource concentration hypothesis, 204
wheat, 204
Spalangia endius,
host age, 346
house fly, 346
parasitization site, 346
parasitoid wasp, 346
pteromalid, 346
Spalangia spp.,
Diapriidae spp., 107
Musca domestica, 107
Pachycrepoideus vindemiae, 107
Stomoxys calcitrans, 107
Trichomalopsis sp., 107
spatial distribution, 280, 517
Callirhytis cornigera, 280
Cynipidae, 280
density-dependent parasitism, 280
diffusion, 517
galls, 280
simulation model, 517
Tripleurospermum inodorum, 517
Triticum aestivum, 517
spatial interpolation,
Cotesia flavipes, 869
parasitoid, 869
spatial-temporal population dynamics, 869
validation, 869
spatial model,
Alphitobius diaperinus, 556
broiler production, 556
geographic information system, 556
lesser mealworm, 556
spatial pattern,
entomopathogenic nematode, 254
population dynamics, 254
spatial-temporal population dynamics,
Cotesia flavipes, 869
parasitoid, 869
spatial interpolation, 869
validation, 869
spider mites,
aerial take-off, 1112
diffusion coefficient, 1112
infochemicals, 1112
walking speed, 1112
spirea aphid,
host suitability, 44
life history, 44
Spissistilus festinus,
Anticarsia gemmatalis, 76
cultural control, 76
Glycine max, 76
Pseudoplusia includens, 76
stink bugs, 76
Spodoptera exigua,
beet armyworm, 511
cotton, 511
egg mass, 511
oviposition, 511
pigweed, 511
Spodoptera littoralis,
aflatoxins, 817
mycotoxins, 817
protein metabolism, 817
spotted knapweed,
Agapeta zoegana, 589
biological control, 589
Centaurea maculosa, 589
Cyphocleonus achates, 589
larval sampling, 589
stage emergence models,
Carposina sasakii, 298
development, 298
starch,
Aedes sollicitans, 388
control, 388
mosquito, 388
xanthan gum, 388
sticky traps,
population monitoring, 663
sex pheromone, 663
Synanthedon tipuliformis, 663
stink bugs,
Anticarsia gemmatalis, 76
cultural control, 76
Glycine max, 76
Pseudoplusia includens, 76
Spissistilus festinus, 76
Stomoxys calcitrans, 107, 147, 982
biological control, 147, 982
Diapriidae spp., 107
life history, 147, 982
Musca domestica, 107, 147, 982
Muscidifurax raptorellus, 982
Muscidifurax zaraptor, 147
Pachycrepoideus vindemiae, 107
Spalangia spp., 107
temperature, 147, 982
Trichomalopsis sp., 107
stored products,
biological control, 617
Oryzaephilus surinamensis, 617
Plodia interpunctella, 617
Rhyzopertha dominica, 617
Tribolium castaneum, 617
sublethal effect,
Choristoneura occidentalis, 136
mortality, 136
multicapsid nucleopolyhedrovirus, 136
nucleopolyhedrovirus, 136
sublethal effects,
Bacillus thuringiensis, 1174
Compsilura, 1174

Lymantria dispar, 1174
subterranean herbivory,
Asclepias, 333
herbivory, 333
Tetraopes, 333
subterranean termites, 673, 832
foraging, 673, 832
landscape mulch, 832
landscapes, 673
resource partitioning, 673
Reticulitermes, 832
Reticulitermes flavipes, 673
Reticulitermes hageni, 673
suction trap,
Aphididae, 371
habitat selection, 371
migratory flight, 371
sugarcane,
arthropods, 1017
cover crops, 1017
Solenopsis invicta, 1017
supercooling,
Eurosta solidaginis, 12
freeze tolerance, 12
over-wintering, 12
water balance, 12
surface airflow,
aggregation sites, 700
dispersal, 700
mate seeking, 700
Ostrinia nubilalis, 700
refuge, 700
surface area,
body water content, 631
cuticular permeability, 631
Dryas iulia, 631
Heliconius, 631
survival, 55, 855
Aedes aegypti, 855
Aedes albopictus, 855
Bemisia tabaci, 55
cantaloupe, 55
cotton, 55
development, 55
fecundity, 55
global warming, 855
larval development, 855
seasonal environment, 855
survivorship,
dry bean, 235
Epilachna varivestis, 235
life tables, 235
susceptibility,
Gracillariidae, 736
Micrurapteryx salicifoliella, 736
Salix, 736
sweet corn,
Ostrinia nubilalis, 274
oviposition, 274
Zea mays, 274
Synanthedon tipuliformis,
population monitoring, 663
sex pheromone, 663
sticky traps, 663
synthetic bait,
component ratio, 1006
pheromone quantity, 1006
Rhyacionia frustrana, 1006
sex pheromone, 1006

T

Tachinidae,
Brachycera, 903
Microtrombidiidae, 903
parasitization, 903
Therevidae, 903
tarnished plant bug,
competition, 1158
in-host compatibility, 1158
Lygus lineolaris, 1158
parasitism, 1158
Peristenus, 1158
Telenomus,
efficient natural control, 37
egg parasitism, 37
Geometridae, 37
Lambdina fiscellaria, 37
outbreak collapse, 37
Temnochila chlorodia,
bark beetles, 993
Enoclerus sphegeus, 993
kairomone, 993
Pityogenes carinulatus, 993
Thanasimus undatulus, 993
temperature, 23, 147, 877, 919, 982, 1090
Adelgidae, 877
biological control, 147, 982
degree-days, 23
density, 919
development, 23
diapause, 877, 1090
egg galleries, 919
flight, 23
forest structure, 919
horn fly, 1090
individual quality, 919
Ips pini, 919
life history, 147, 982
Musca domestica, 147, 982
Muscidifurax raptorellus, 982
Muscidifurax zaraptor, 147
overwintering survival, 1090
phenological model, 23
photoperiod, 877
population dynamics, 1090
Sesamia nonagrioides, 23
Stomoxys calcitrans, 147, 982
Tsuga spp. aestivation, 877
temperature response,
calorespirometry, 811
convergent lady beetle, 811
Hippodamia convergens, 811
isothermal calorimetry, 811
respiration, 811
temperature-dependent growth,
degree-day models, 692
developmental models, 692
grape berry moth, 692
temperatures,
aphids, 1127
corn, 1127
corn aphids, 1127
development, 1127
fecundity, 1127
termite,
bacteria, 443
bio-control, 443
Coptotermes formosanus, 443
fungus, 443
Serratia marcescens, 443
termite tunneling,
Coptotermes formosanus, 545
Reticulitermes flavipes, 545
tunneling geometry, 545
Tetranychus urticae,
foraging behavior, 1103
predaceous mites, 1103
predator specialization, 1103
prey switching, 1103
Tetraopes,
Asclepias, 333
herbivory, 333
subterranean herbivory, 333
Thanasimus undatulus,
bark beetles, 993
Enoclerus sphegeus, 993
kairomone, 993
Pityogenes carinulatus, 993
Temnochila chlorodia, 993
Therevidae,
Brachycera, 903
Microtrombidiidae, 903
parasitization, 903
Tachinidae, 903
thermal thresholds,
development, 1082
dry cured-ham, 1082
models, 1082
mold mite, 1082
Tyrophagus putrescentiae, 1082
thermoperiod,
fat content, 166
overwintering, 166
photoperiod, 166
Reticulitermes flavipes, 166
water content, 166
thermoregulation,
feeding behavior, 172
herbivory, 172
Japanese beetle, 172
Popillia japonica, 172

thresholds,
avocado, 932
scarring on avocado fruit, 932
Scirtothrips perseae, 932
thrips numbers, 932
thrips numbers,
avocado, 932
scarring on avocado fruit, 932
Scirtothrips perseae, 932
thresholds, 932
ticks,
Amblyomma americanum, 568
distribution, 568
habitats, 568
Ixodes scapularis, 568
tomato yellow leaf curl virus,
flight behavior, 394
pest management, 394
ultraviolet light, 394
whiteflies, 394
tortrix,
Cydia trasias, 7
mating disruption, 7
sex pheromone, 7
Toxoptera citricida,
alatae, 1047
apterae, 1047
density, 1047
development, 1047
trail pheromone,
chemical communication, 457
Coptotermes formosanus, 457
Reticulitermes flavipes, 457
transgenic,
Bacillus thuringiensis, 495
Bt-corn, 495
Danaus plexippus, 495
monarch butterfly, 495
nontarget effects, 495
transgenic classical resistance,
Bacillus thuringiensis tenebrionis, 524
Colorado potato beetle, 524
plant–insect interactions, 524
potato, 524
potato aphid, 524
transgenic corn,
Bacillus thuringiensis, 409
insect resistance management, 409
Ostrinia nubilalis, 409
pollen, 409
Zea mays, 409
trap,
alfalfa looper, 667
attractant, 667
Autographa californica 667
phenylacetaldehyde, 667
trapping, 64, 656
acetic acid, 656
attractant, 656
boll weevil, 64
cultural management, 64
Lacanobia subjuncta, 656
Mamestra configurata, 656
overwintering, 64
pheromone synergist, 64
Xestia c-nigrum, 656
traps, 82, 318
dispersal, 318
flies, 82
mite management, 318
Phyllocoptruta oleivora, 318
Rhagoletis cingulata, 82
Rhagoletis fausta, 82
Trialeurodes vaporariorum,
Bemisia tabaci, 89
intercropping, 89
polyculture, 89
vector management, 89
Tribolium castaneum,
biological control, 617
Oryzaephilus surinamensis, 617
Plodia interpunctella, 617
Rhyzopertha dominica, 617
stored products, 617
Trichomalopsis sp.,
Diapriidae spp., 107
Musca domestica, 107
Pachycrepoideus vindemiae, 107
Spalangia spp., 107
Stomoxys calcitrans, 107
Tripleurospermum inodorum,
diffusion, 517
simulation model, 517
spatial distribution, 517
Triticum aestivum, 517
Triticum aestivum, 225, 517
agroecosystems, 225
boundaries, 225
diffusion, 517
dispersal, 225
multivariate analysis, 225
predators, 225
simulation model, 517
spatial distribution, 517
Tripleurospermum inodorum, 517
tritrophic interactions, 947, 964
Acyrthosiphon pisum, 964
Brassicaceae, 947
Coccinellidae, 964
Coleomegilla maculata, 964
fatty acids, 964
Hippodamia convergens, 964
lady beetle, 947
Myzus persicae, 947
trophic cascade,
antipredator behavior, 327
Diabrotica undecimpunctata howardi, 327
predation, 327
sexual differences, 327
Tsuga spp. aestivation,
Adelgidae, 877
diapause, 877
photoperiod, 877
temperature, 877
tunneling geometry,
Coptotermes formosanus, 545
Reticulitermes flavipes, 545
termite tunneling, 545
turfgrass,
ants, 425
Cyclocephala lurida, 425
Popillia japonica, 425
predators, 425
Solenopsis molesta, 425
Tyrophagus putrescentiae,
development, 1082
dry cured-ham, 1082
models, 1082
mold mite, 1082
thermal thresholds, 1082

U

ultraviolet light,
flight behavior, 394
pest management, 394
tomato yellow leaf curl virus, 394
whiteflies, 394
United States,
biogeographical range, 645
distribution, 645
modeling, 645
quarantine, 645
Solenopsis invicta, 645
Urbanus proteus,
Agraulis vanillae, 1052
Eurema lisa, 1052
Junonia coenia, 1052
phenology, 1052
Phoebis sennae, 1052

V

validation,
Cotesia flavipes, 869
parasitoid, 869
spatial interpolation, 869
spatial-temporal population dynamics, 869
vector management,
Bemisia tabaci, 89
intercropping, 89
polyculture, 89
Trialeurodes vaporariorum, 89
vertical distribution of insect fauna,
clearcut, 400
forest management, 400
insect sampling, 400
selection, 400
shelterwood, 400

Vespidae,
Attractants, 157
food volatiles, 157
isobutanol, 157
Vespula germanica, 157
yellowjackets, 157
Vespula germanica,
Attractants, 157
food volatiles, 157
isobutanol, 157
Vespidae, 157
yellowjackets, 157

W

walking,
colonization, 306
flight, 306
impact traps, 306
Leptinotarsa decemlineata, 306
plastic-lined trenches, 306
walking speed,
aerial take-off, 1112
diffusion coefficient, 1112
infochemicals, 1112
Spider mites, 1112
warning coloration,
aggregation, 51
ambush predator, 51
chemical defense, 51
frogs, 51
water balance,
Eurosta solidaginis, 12
freeze tolerance, 12
over-wintering, 12
supercooling, 12
water content,
fat content, 166
overwintering, 166
photoperiod, 166
Reticulitermes flavipes, 166
thermoperiod, 166
water stress,
Diprion pini, 1
foliage dry matter, 1
hatching, 1
larval survival, 1
Pinus sylvestris, 1
web spiders,
ambushers, 1021
diversity, 1021
hunting spiders, 1021
predatory arthropods, 1021
sampling methods, 1021
weeds,
Bacillus thuringiensis, 728
corn, 728
larval survival, 728
Ostrinia nubilalis, 728
resistance management, 728
weevil trapping,
aquatic barrier trap, 770
aquatic intercept trap, 770
Lissorhoptrus oryzophilus, 770
rice integrated pest management, 770
rice water weevil, 770
western spruce budworm,
Acleris gloverana, 883
age-specific survival ratios, 883
dispersal, 883
Douglas-fir tussock moth, 883
population dynamics, 883
wheat,
cultural control, 204
Empoasca fabae, 204
intercropping, 204
resource concentration hypothesis, 204
soybean, 204
whiteflies,
flight behavior, 394
pest management, 394
tomato yellow leaf curl virus, 394
ultraviolet light, 394
winter mortality,
cold hardiness, 181
invading species, 181
range limit, 181
southwestern corn borer, 181
within-plant variation,
developmental resistance, 1144
Eucalyptus globulus, 1144
herbivory, 1144
heteroblasty, 1144
phase change, 1144

X

xanthan gum,
Aedes sollicitans, 388
control, 388
mosquito, 388
starch, 388
Xestia c-nigrum,
acetic acid, 656
attractant, 656
Lacanobia subjuncta, 656
Mamestra configurata, 656
trapping, 656
Xyleborinus saxeseni,
flight patterns, 909
insect-plant interactions, 909
Scolytidae, 909
Xylosandrus crassiusculus, 909
Xylosandrus germanus, 909
Xylosandrus crassiusculus,
flight patterns, 909
insect-plant interactions, 909
Scolytidae, 909
Xyleborinus saxeseni, 909
Xylosandrus germanus, 909
Xylosandrus germanus,
flight patterns, 909
insect-plant interactions, 909
Scolytidae, 909
Xyleborinus saxeseni, 909
Xylosandrus crassiusculus, 909

Y

yellowjackets,
Attractants, 157
food volatiles, 157
isobutanol, 157
Vespidae, 157
Vespula germanica, 157

Z

Zea mays, 274, 409
Bacillus thuringiensis, 409
insect resistance management, 409
Ostrinia nubilalis, 274, 409
oviposition, 274
pollen, 409
sweet corn, 274
transgenic corn, 409

Information for Contributors

Environmental Entomology is published in February, April, June, August, October, and December. Contributors report on the interaction of insects with the biological, chemical, and physical aspects of their environment. Articles are divided into categories by subject matter: physiological and chemical ecology (abiotic effects, pheromones, effects of miscellaneous pollutants), community/ecosystem ecology (trophic-level studies, associations), population ecology (mating, reproduction, movement, behavior, parasitism, predation, microbial ecology, insect–plant relations), pest management and sampling (integrated pest management, sampling, distribution), and biological control (mass release, microbial control, augmentation, classical biological control).

All manuscripts are reviewed by scientists qualified to judge the validity of the research. Acceptance or rejection, however, is the decision of the subject editor. Acceptance of manuscripts is based solely on their scientific merit. Appeal of a rejected manuscript should be made to the Editorial Board, via the Director of Publications. Accepted manuscripts are published approximately in the order they are received.

In addition to research papers, *Environmental Entomology* publishes *Letters to the Editor*, interpretive articles in a *Forum* section, *Book Reviews*, and articles that describe new entomological techniques. The same guidelines apply to these as to research articles.

For more detailed instructions concerning the preparation and submission of manuscripts, consult the *Publishing with ESA Style Guide* published on our web site, http://www.entsoc.org.

Manuscript Preparation. Type all manuscripts double-spaced, with 1-inch margins. Number pages consecutively, beginning with the title page. Begin each of the following on a separate page and arrange in the following order: title page, abstract and key words (three to six words), text, acknowledgments, references cited, footnotes, tables, and figure legends. Submit an original and two copies of the manuscript and artwork to:

Environmental Entomology
Entomological Society of America
9301 Annapolis Road
Lanham, MD 20706

Do not submit a new manuscript directly to the subject editor.

Title Page. In the upper right corner, type the complete name, address, and telephone number, fax number, and email of the author who is to receive and approve page proofs. In the upper left corner, type *Environmental Entomology*, the section of the journal in which the manuscript should appear, and the running head. Use a brief and informative title that includes order and family names in parentheses. Four lines beneath the title, type in capital letters the complete names of all authors. Four lines beneath the authors' names, type the department, institution, city, state, zip code, and country (other than the United States) of the institution where the research was done. Use numbered footnotes for authors not affiliated with the research institution, and give their addresses or affiliations on the footnote page.

Abstract. Abstract should be an informative digest of the significant contents and the main conclusions of the research.

References Cited. Citations in the text should be by last name of the author and date and refer to a reference listed alphabetically by author in References Cited. For further formatting instructions, see our *Publishing with ESA Style Guide* located on our Web site (http://www.entsoc.org).

Tables. Keep tables to a minimum. Type all parts of each table double-spaced, with each table starting a new page.

Illustrations. At the time of manuscript submission, submit original drawings. Photographs must be on glossy paper. Separate parts of the same figure must be mounted together on illustration board and arranged to use space efficiently. Authors should indicate a preference for one- or two-column reproduction, and they are asked to account for the length of the caption and for proportionated reduction or enlargement in the choice of lettering size, type style, and the format of the artwork itself. Final (reduced) lettering size should be 10 points. Authors are urged to contact the staff of the publications office with questions concerning the preparation of artwork. Each piece of artwork should be labeled with the corresponding author's name, address, and telephone number; an abbreviated title of the paper; and the name of the journal to which the article has been submitted. Original artwork for rejected papers is returned by the subject editor; all other artwork is returned to the corresponding author upon request.

Style Conventions. When first mentioned, a plant or animal should include the full scientific name and the author of the zoological name unabbreviated—except for Linnaeus (L.) and Fabricius (F.). Use only common names approved by the Entomological Society of America (Common Names of Insects & Related Organisms 1997). Otherwise, the manuscript should conform to the guidelines established by the latest edition of the *Council of Biology Editors Style Manual.*

Proofs and Reprints. Page proofs and reprint order forms are sent to authors after manuscripts are typeset. All changes in proof, except for printer and editorial errors and answers to copy editor queries addressing style or format, will be charged to the author.

Editorial Charges. Editorial review charges are assessed on all accepted manuscripts to cover some of the expenses of publishing. Contact the publications office at (301) 731-4535 or the ESA web site (http://www.entsoc.org) for current rates. Reprints (printed and electronic) are ordered at the time page proofs are returned to ESA; see the reprint order form for rates. Author alterations are billed at $6 per alteration (alterations to figures are billed at a higher rate). Member authors who are retired and not affiliated with any institution can publish five pages gratis each year. Member authors who are employed but who have no grants or funds for publication can request a partial waiver of editorial charges. Such requests must be made at the time of manuscript submission. For a full explanation, see our publications policies on the ESA web site (http://www.entsoc.org).

THIS PUBLICATION IS PRINTED ON ACID-FREE, RECYCLED PAPER.